






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<div>(51) International Patent Classification ⁶ : A61K 31/495, 31/535</div>	<div>A1</div>	<div>(11) International Publication Number: WO 99/13877</div> <div>(43) International Publication Date: 25 March 1999 (25.03.99)</div>
<div>(21) International Application Number: PCT/SE98/01601</div> <div>(22) International Filing Date: 9 September 1998 (09.09.98)</div> <div>(30) Priority Data: 9703375-7 18 September 1997 (18.09.97) SE</div> <div>(71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE).</div> <div>(72) Inventors; and</div> <div>(75) Inventors/Applicants (for US only): BERG, Stefan [SE/SE]; Astra Arcus AB, S-151 85 Södertälje (SE). ROSS, Svante [SE/SE]; Astra Arcus AB, S-151 85 Södertälje (SE). THORBERG, Seth-Olov [SE/SE]; Astra Arcus AB, S-151 85 Södertälje (SE).</div> <div>(74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).</div>		<div>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</div> <div>Published</div> <div>With international search report.</div> <div>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</div>

(54) Title: A COMBINATION OF A 5-HT REUPTAKE INHIBITOR AND A h5-HT_{1B} ANTAGONIST OR PARTIAL AGONIST

(57) Abstract

The invention relates to a combination of a first component (a) which is a 5-HT reuptake inhibitor and a second component (b) which is selective h5-HT_{1B} antagonist or partial agonist having formula (I) wherein X is CH₂, O; Y is CONH, NHCO; R₁ is H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl; R₂ is H, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen; R₃ is (1), (2), (3), CF₃; R₄ and R₅ independently are H, or C₁-C₄ alkyl, as racemate, R-enantiomer or S-enantiomer, and said components (a) and (b) being in the form of free bases, solvents or pharmaceutically acceptable salts thereof, the preparation thereof, pharmaceutical formulations containing said combinations, use of and method of treatment of affective disorders such as depression, anxiety and OCD with said combination as well as kit containing said combination.



● Compound A 12 µmol/kg +
Compound B 20 µmol/kg
 ■ Compound A 12 µmol/kg +
Compound C 2 ml/kg
 ▼ Compound A 12 µmol/kg +
Compound D 23 µmol/kg
 ◆ Compound A 12 µmol/kg +
Compound E 20 µmol/kg

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A combination of a 5-HT reuptake inhibitor and a h5-HT_{1B} antagonist or partial agonist

5 **Field of the Invention**

The present invention relates to a product which comprises a **combination of a 5-HT** reuptake inhibitor (SSRI) and a selective h5-HT_{1B} receptor antagonist or partial agonist , more specifically a piperidyl- or piperazinyl-substituted 1,2,3,4-tetrahydronaphthalene or -
10 3,4-dihydro-2*H*-1-benzopyran derivative whereby each components are in the form of the free bases, solvates or pharmaceutically acceptable salts thereof. The present invention also relates to a process for the preparation of the inventive combination, pharmaceutical formulation comprising said combination and to the use of said combination by either concomitant administration or separate administration as an improvement of the treatment
15 of affective disorders such as depression, anxiety, obsessive compulsive disorder (OCD), *etc.*

Background of the Invention

20

Today, it is generally considered that antidepressants, including Selective Serotonin (5-HT) Reuptake Inhibitors (SSRIs), take 2-4 weeks to reach their full clinical effect. In contrast, the side effects occur immediately. Thus, the slow onset of action of antidepressants leads to a vulnerable period for patients in which they experience the side effects, but not the
25 therapeutic effects of drugs. There is often a heavy burden on the treating physician to persuade the patient to continue with the treatment during this period. Furthermore, in suicidal patients, as the onset of action is gradual, they may regain initiative without experiencing full reversal symptoms, leading to a window of risk for suicide and a frequent requirement for hospitalisation. An antidepressant with fast onset of action would not only
30 be beneficial due to the faster symptom reduction, but would also be more acceptable to

patients and physicians and reduce the need for and duration of hospitalisation. The same long period to reach full clinical effect has been shown in the treatment of other affective disorders such as anxiety and OCD.

5 **Summary of the Invention**

The present invention is directed, in part, to a combination of SSRI and a 5HT_{1B} receptor antagonist.


10 Advantageously, treatment with the combination results in an increased synaptic concentration of 5HT. SSRIs decrease the release of 5-HT in 5-HT neurons via a negative feed-back reaction in part mediated by terminal h5-HT_{1B} receptors. By inhibiting the terminal h5-HT_{1B} autoreceptors, the selective receptor antagonists counteract the decrease in 5-HT release caused by 5-HT reuptake inhibitors. This indicates that a selective
15 blockade of terminal autoreceptors by a h5-HT_{1B} antagonist may have a clinical potential to improve the efficacy of 5-HT reuptake inhibitors (SSRIs) and offer a new rational for rapid onset of affective actions, for instance the antidepressant actions.

The combination

20

Thus, by combining a first component (a) which is a 5-HT reuptake inhibitor with a second component (b) which is a selective h5-HT_{1B} antagonist or partial agonist having the formula I

25




wherein X is CH₂, O;

Y is CONH, NHCO;

R₁ is H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl;

R₂ is H, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen;

R₃ is



R₄ and R₅ independently are H or C₁-C₄ alkyl,

as racemate, *R*-enantiomer or *S*-enantiomer, and said components (a) and (b) being in the form of free bases, solvates, preferably hydrates or pharmaceutically acceptable salts thereof, a faster onset of action will occur and consequently, a more efficacious treatment of the patients.

In other preferred embodiments of the second component (b) are those compounds of

formula I wherein X is CH₂, and of those compounds, compounds wherein Y is NHCO,

and of those compounds, compounds wherein R₃ is morpholino. Compounds wherein R₁

is hydrogen, methyl or ethyl and wherein R₂ is hydrogen, methyl, ethyl, methoxy or bromo are preferred.

5 Preferred compounds having the formula I are:

(*R*)-*N*-[8-(Piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[8-(4-Ethylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

10 (*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[5-Ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

15 (*R*)-*N*-[5-Ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-(4-morpholinocarbonyl)benzamide;

(*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinocarbonylbenzamide;

(*R*)-*N*-[5-Bromo-8-(piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

20 *N*-[5-Bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[5-Bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-trifluoromethylbenzamide;

25 (*R*)-*N*-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

N-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

(*R*)-*N*-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

(*S*)-*N*-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

(*R*)- *N*-(Morpholinocarbonylphenyl)-8-(4-methylpiperazin-1-yl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

5 (*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-morpholinobenzamide;

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(4-piperidon-1-yl)benzamide;

10 (*S*)-*N*-[8-Methyl-5-(4-methyl-piperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(dimethylaminocarbonyl)benzamide;

N-[4-(4-Morpholinyl)phenyl]-8-methoxy-5-(4-methyl-piperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-carboxamide.

Particularly preferred compounds are (*R*)-*N*-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide, (*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide and (*R*)-*N*-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide.

20 The compounds of formula I as (*R*)-enantiomers, (*S*)-enantiomers or racemates may exist in the form of a free base or a pharmaceutically acceptable salt or hydrate thereof.

In the present context C₁-C₆ alkyl may be straight or branched. C₁-C₆ alkyl may be methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, t-pentyl, neo-pentyl, n-hexyl or i-hexyl. Preferred is C₁-C₄ alkyl and especially preferred are
25 methyl and ethyl.

In the present context C₁-C₄ alkyl may be straight or branched. C₁-C₄ alkyl may be methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl or t-butyl. Methyl and ethyl are preferred.

In the present context C₃-C₆ cycloalkyl may be cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

In the present context C₁-C₆ alkoxy may be straight or branched. C₁-C₆ alkoxy may be
5 methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, i-butoxy, s-butoxy, t-butoxy, n-pentyloxy, i-pentyloxy, t-pentyloxy, neo-pentyloxy, n-hexyloxy or i-hexyloxy. Preferred is C₁-C₄ alkoxy and especially preferred is methoxy.

10 In the present context halogen may be fluoro, chloro, bromo or iodo wherein bromo is preferred.

Suitable known 5-HT reuptake inhibitors (SSRIs) to be used are norzimeldine, fluoxetine, paroxetine, citalopram, clomipramine, sertraline, fluvoxamine, alaproclate, preferably citalopram or paroxetine, but component (a) in the combination according to the invention
15 is not limited only to these SSRIs.

The combination according to the present invention may exist in one pharmaceutical formulation comprising both the active first component (a) and the active second component (b) or in two different pharmaceutical formulations, one for the active first
20 component (a) and one for the active second component (b). The pharmaceutical formulation may be in the form of tablets or capsules, powders, mixtures, solutions or other suitable pharmaceutical formulation forms.

The combination of the present invention can be prepared by incorporating a 5-HT
25 reuptake inhibitor is incorporated into the same formulation as a selective h5-HT_{1B} antagonist as defined above, e.g. by mixing in a conventional way.

The present invention also includes a method of improving the onset of therapeutic action by concomitant administration of a combination of a first component (a) which is a 5-HT

reuptake inhibitor and a second component (b) which is selective h5-HT_{1B} antagonist as defined above.

5 A further embodiment of the present invention is a kit containing the combination of a first component (a) which is a 5-HT reuptake inhibitor and a second component (b) which is selective h5-HT_{1B} antagonist as defined above. The kit may include an instruction for use.

Pharmaceutical formulations

10 According to the present invention the compounds in the combination will normally be administered orally, rectally or by injection, in the form of pharmaceutical formulations comprising the active ingredient either as a free base, solvates e.g. hydrates or a pharmaceutically acceptable non-toxic acid addition salt, e.g. the hydrochloride, hydrobromide, lactate, acetate, phosphate, sulfate, sulfamate, citrate, tartrate, oxalate and
15 the like in a pharmaceutically acceptable dosage form. The dosage form may be a solid, semisolid or liquid formulation. Usually the active substances will constitute between 0.1 and 99% by weight of the formulation, more specifically between 0.5 and 20% by weight for formulations intended for injection and between 0.2 and 50% by weight for formulations suitable for oral administration.

20 The pharmaceutical formulation comprises the active ingredients optionally in association with adjuvants, diluents, excipients and/or inert carriers.

To produce pharmaceutical formulations of the combination of the invention in the form of dosage units for oral application, the selected compounds may be mixed with a solid excipient, e.g. lactose, saccharose, sorbitol, mannitol, starches such as potato starch, corn
25 starch or amylopectin, cellulose derivatives, a binder such as gelatine or polyvinylpyrrolidone, disintegrants e.g. sodium starch glycolate, cross-linked PVP, cross-carmellose sodium and a lubricant such as magnesium stearate, calcium stearate, polyethylene glycol, waxes, paraffin, and the like, and then compressed into tablets. If coated tablets are required, the cores, prepared as described above, may be coated with a
30 concentrated sugar solution which may contain e.g. gum arabic, gelatine, talcum, titanium

dioxide, and the like. Alternatively, the tablets can be coated with a polymer known to the man skilled in the art, wherein the polymer is dissolved in a readily volatile organic solvent or mixture of organic solvents. Dyestuffs may be added to these coatings in order to readily distinguish between tablets containing different active substances or different amounts of the active compounds.

For the formulation of soft gelatine capsules, the active substances may be admixed with e.g. a vegetable oil or poly-ethylene glycol. Hard gelatine capsules may contain granules of the active substances using either the above mentioned excipients for tablets e.g. lactose, saccharose, sorbitol, mannitol, starches (e.g. potato starch, corn starch or amylopectin), cellulose derivatives or gelatine. Also liquids or semisolids of the drug can be filled into hard gelatine capsules.

Dosage units for rectal application can be solutions or suspensions or can be prepared in the form of suppositories comprising the active substances in a mixture with a neutral fatty base, or gelatine rectal capsules comprising the active substances in admixture with vegetable oil or paraffin oil. Liquid formulations for oral application may be in the form of syrups or suspensions, for example solutions containing from about 0.2% to about 20% by weight of the active substances herein described, the balance being sugar and mixture of ethanol, water, glycerol and propylene glycol. Optionally such liquid formulations may contain colouring agents, flavouring agents, saccharine and carboxymethyl-cellulose as a thickening agent or other excipients known to a person skilled in the art.

Solutions for parenteral applications by injection can be prepared in an aqueous solution of a water-soluble pharmaceutically acceptable salt of the active substances, preferably in a concentration of from about 0.5% to about 10% by weight. These solutions may also contain stabilizing agents and/or buffering agents and may conveniently be provided in various dosage unit ampoules.

Suitable daily doses of the active compounds in the combination of the invention in therapeutical treatment of humans are about 0.01-100 mg/kg bodyweight on peroral

administration and 0.001-100 mg/kg bodyweight on parenteral administration. The daily doses of the active h5-HT_{1B} antagonist may very well differ from the daily doses of the SSRIs but the doses can also be the same for both of the active compounds.

5

Medical and Pharmaceutical Use


In a further aspect the present invention provides the use of the combination of a first component (a) which is a 5-HT reuptake inhibitor with a second component (b) which is a
10 selective h5-HT_{1B} antagonist or partial agonist, preferably an antagonist, having the formula I as defined herein, and the use in the treatment of 5-hydroxytryptamine mediated disorders, such as affective disorders. Examples of affective disorders are disorders in the CNS such as mood disorders (depression, major depressive episodes, dysthymia, seasonal affective disorder, depressive phases of bipolar disorder), anxiety disorders (obsessive
15 compulsive disorder, panic disorder with/without agoraphobia, social phobia, specific phobia, generalized anxiety disorder, posttraumatic stress disorder), personality disorders (disorders of impulse control, trichotellomania). Other disorders in CNS such as obesity, anorexia, bulimia, premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders (age
20 associated memory impairment, presenile and senile dementia), pathological aggression, schizophrenia, endocrine disorders (e g hyperprolactinaemia), stroke, dyskinesia, Parkinson's disease, thermoregulation, pain, hypertension may be treated with the combination described herein, too. Examples of other hydroxytryptamine mediated disorders are urinary incontinence, vasospasm and growth control of tumors (e g lung
25 carcinoma) and it may be possible to treat those with the combination described herein as well.

30

Method of Preparation of Intermediates.


1. In the case where Y is NHCO and X is CH₂ or O.

- 5 (i) Benzylation of the compound of the formula **II**, either as a racemate or as an enantiomer,




- 10 to obtain a compound of formula **III** may be carried out by reaction with a suitable benzylation agent e.g. a benzyl halide such as benzyl bromide or benzyl chloride or an activated alcohol e.g. benzyl mesylate or benzyl tosylate. The reaction may be carried out using a salt or the base of compound **II** in a suitable solvent e.g. *N,N*-dimethylformamide, acetone or acetonitrile with a suitable base e.g. NaOH, NaHCO₃, K₂CO₃ or a
- 15 trialkylamine such as triethylamine at a temperature within the range of +20 °C to +150 °C. The presence of a suitable catalyst e.g. potassium iodide or sodium iodide, may increase the speed of the reaction. The nitrogen in compound **II** may also be protected by reductive alkylation with an arylaldehyde in the presence of a reductive agent such as sodium cyanoborohydride, sodium borohydride or catalytically with H₂ and a suitable catalyst
- 20 containing palladium, platinum, rhodium or nickel in a suitable solvent e.g. tetrahydrofuran, dioxane, methanol or ethanol. A proton donor such as p-toluenesulfonic acid can be used to catalyze the formation of the imine/enamine, and adjustment of pH to slightly acidic by an appropriate acid such as acetic acid may speed up the reaction, resulting in compound **III**.

(ii) Demethylation of the compound of formula III




5 to obtain a compound of formula IV may be carried out by treating the compound with an acidic reagent such as aqueous HBr, HI, HBr/CH₃COOH, BBr₃, AlCl₃, pyridine-HCl or with a basic nucleophilic reagent such as CH₃C₆H₄S⁻ or C₂H₅S⁻ in a suitable solvent. Suitable solvents may be methylene chloride or chloroform and the reaction may occur
10 between -78 °C and +60 °C.

(iii) Conversion of the compound of formula IV to a compound of formula V



15 may be carried out by the reaction with a compound of formula VI




20 where L stands for a leaving group, e.g. a halogen such as chlorine, bromine or iodine or an alkane- or arenesulfonyloxy group such as a p-toluenesulfonyloxy group and Ra and Rb are hydrogen or a lower alkyl group e.g. methyl. The process may be carried out with a salt of the compound of formula IV obtained by reaction with a base such as K₂CO₃, Na₂CO₃,

KOH, NaOH, BuLi or NaH. The reaction may be conducted in a suitable solvent e.g. an aprotic solvent such as dioxane, *N,N*-dimethylformamide, tetrahydrofuran, toluene, benzene or petroleum ether and the reaction may occur between +20 °C and +150 °C.

5


(iv) Rearrangement of a compound of formula **V** to a compound of formula **VII**



10 may be carried out in a suitable solvent e.g. aprotic solvent such as *N,N*-dimethylformamide, dioxane, 1,1,3,3-tetramethylurea, tetrahydrofuran or hexamethylphosphoric triamide with a suitable base e.g. K_2CO_3 , KOH, potassium *tert*-butoxide or NaH at a temperature within the range of +20 °C to +150 °C. The presence of a cosolvent such as 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone or
 15 hexamethylphosphoric triamide in appropriate concentration in the solvent may increase the speed of the reaction.


(v) Hydrolysis of a compound of formula **VII** to a compound **VIII** may be carried out under acidic conditions using acids such as H_2SO_4 , HCl or HBr in a suitable solvent e.g.
 20 H_2O , ethanol, methanol or mixtures thereof and the reaction may occur between +20 °C and +100 °C or under basic conditions using bases such as NaOH or KOH in a suitable solvent e.g. H_2O , ethanol, methanol or mixtures thereof and the reaction may occur between +20 °C and +100 °C.

25 (vi) Conversion of compound of formula **VIII** to a compound of formula **IX**




may be carried out by

- 5 a) reaction with a compound of formula X



- 10 where R_1 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl. The process may be carried out in a suitable solvent e.g. an aprotic/anhydrous solvent such as tetrahydrofuran or *N,N*-dimethylformamide in the presence of coupling reagent such as *N,N'*-carbonyldiimidazole and the reaction may occur between +20 °C and +130 °C. The reaction is followed by the reduction of the imide with a suitable reducing agent e.g. $LiAlH_4$ in a suitable solvent e.g.
- 15 diethyl ether or tetrahydrofuran at a temperature between +20 °C and reflux, or

- b) by reaction with a compound of formula XI




where L stands for a leaving group, e.g. a halogen such as chlorine or bromine or an alkane- or arenesulfonyloxy group such as p-toluenesulfonyloxy group and R_1 is H, C_1 - C_6 -alkyl or C_3 - C_6 cycloalkyl. The process may be carried out in a suitable solvent such as ethanol, butanol, *N,N*-dimethylformamide, acetonitrile or a mixture of water and
5 acetonitrile with a suitable base e.g. K_2CO_3 , $NaHCO_3$ or KOH and the reaction may occur between +20 °C and +150 °C.

Conversion of a compound of formula IX, wherein R_1 is hydrogen, to an alkylated compound of formula IX, where R_1 is C_1 - C_6 alkyl, may be carried out by using a suitable
10 alkylation reagent such as R_1 -L, where L is a suitable leaving group e.g. a halogen such as chlorine, bromine or iodine or an alkane- or arenesulfonyloxy group such as a p-toluenesulfonyloxy group and R_1 is C_1 - C_6 alkyl. The reaction may be carried out in a suitable solvent such as *N,N*-dimethylformamide, acetone, acetonitrile or tetrahydrofuran with a suitable base such as K_2CO_3 , $NaHCO_3$, NaOH or a trialkylamine such as
15 triethylamine. The reaction may be conducted at a temperature between +20 °C and + 120 °C, or


conversion of a compound of formula IX, where R_1 is hydrogen, to an alkylated compound of formula IX, where R_1 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, may be carried out by
20 reductive alkylation with a compound R_1 -CHO, where R_1 is hydrogen or C_1 - C_5 alkyl, or with a C_3 - C_6 cyclic ketone, in the presence of a reductive agent, such as sodium cyanoborohydride, sodium borohydride or catalytically with H_2 and a suitable catalyst containing palladium, platinum, rhodium or nickel in a suitable solvent e.g. tetrahydrofuran, dioxane, methanol or ethanol. A proton donor such as p-toluenesulfonic
25 acid can be used to catalyze the formation of the imine/enamine and adjustment of pH to slightly acidic by an appropriate acid such as acetic acid may speed up the reaction.

(vii) Halogenation of the compound of formula **IX**, where R_1 is hydrogen, C_1 - C_6 -alkyl or C_3 - C_6 -cycloalkyl,



to obtain a compound of formula **XII** may be performed by aromatic electrophilic substitution using a suitable halogenation agent such as Br_2 , Cl_2 , I_2 , ICl , or SO_2Cl_2 . The reaction may be carried out using the salt or the base of the compound **IX** in an appropriate solvent e.g. acetic acid, HCl /ethanol or water with or without a suitable base e.g. alkali metal acetate such as sodium acetate and at a reaction temperature between $-20\text{ }^{\circ}C$ and room temperature.


10




- (viii) Conversion of the compound of formula **XII** to a compound of formula **XIII**, where R_1 is hydrogen, C_1-C_6 alkyl or C_3-C_6 cycloalkyl and R_2 is C_1-C_6 alkyl, may be carried out by a metal-halogen exchange, in a appropriate anhydrous solvent such as tetrahydrofuran or diethyl ether using a suitable alkyl-lithium or metal e.g. butyllithium, lithium or magnesium turnings, followed by treatment with appropriate alkyl halide such as methyl iodide, ethyl bromide or propyl iodide and the reaction may be performed at a reaction temperature within the range of -78°C to room temperature, followed by cleavage of the benzyl groups by hydrogenation over a suitable catalyst containing palladium, rhodium, platina or nickel, in a suitable solvent e.g. acetic acid or ethanol and at a reaction temperature between $+20^\circ\text{C}$ and $+120^\circ\text{C}$, or treatment with other electrophiles such as acetaldehyde or methyl chloroformate and a thereafter following suitable work-up. The reaction may be performed at a reaction temperature within the range of -78°C to room temperature.
- 15 In the case where acetaldehyde is used as electrophile, the above reaction is followed by reduction of the benzyl alcohol and cleavage of the benzyl groups by hydrogenation over a suitable catalyst containing palladium, rhodium, platina or nickel, in a suitable solvent e.g. acetic acid or ethanol and the reaction may occur between $+20^\circ\text{C}$ and $+120^\circ\text{C}$.
- 20 In the case where methyl chloroformate is used as electrophile, the above reaction is followed by reduction of the methyl ester in a suitable solvent such as diethyl ether or tetrahydrofuran with an appropriate reductive agent such as lithium aluminum hydride and the reaction may occur between $+20^\circ\text{C}$ and reflux, followed by cleavage of the benzyl groups and reduction of the benzyl alcohol by hydrogenation over a suitable catalyst
- 25 containing palladium, rhodium, platina or nickel, in a suitable solvent e.g. acetic acid or ethanol and the reaction may occur between $+20^\circ\text{C}$ and $+120^\circ\text{C}$.

When R_1 is hydrogen, the piperazine nitrogen is protected with a suitable protecting group before the lithiation step such as a benzyl group or another protecting group known by a person skilled in the art and then removed by methods known by a person skilled in the art, resulting in the compound of formula **XIII**.

(ix) Conversion of a compound of formula **XIII**, where R_1 is hydrogen, to a compound of formula **XIV**,




where R_c is a suitable protecting group, may be carried out by the protection of the piperazine ring in a suitable solvent e.g. methylene chloride or chloroform with a
 10 appropriate protecting reagent e.g. di-*tert*-butyl dicarbonate with a suitable base e.g. triethylamine or K_2CO_3 and at a temperature between $-20\text{ }^{\circ}\text{C}$ and $+60\text{ }^{\circ}\text{C}$.



(x) Conversion of the compound of formula **IX**, where R_1 is hydrogen, C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, to a compound of formula **XV**, where R_1 is hydrogen, C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl may be carried out by cleavage of the benzyl groups by hydrogenation over a

suitable catalyst containing palladium, rhodium, platina or nickel, in a suitable solvent e.g. acetic acid or ethanol and the reaction may occur between +20 °C and +120 °C.

(xi) Conversion of a compound of formula **IX** where R₁ is hydrogen, to a compound of formula **XVI**




where R_c stands for a suitable protecting group, may be carried out by

a) hydrogenation using a catalyst containing palladium, platinum, rhodium or nickel in a suitable solvent e.g. acetic acid or ethanol at a reaction temperature between +20 °C and +120 °C, or

b) debenzylation in a suitable solvent such as methanol in the presence of ammonium formate and Pd/C at a reaction temperature between +20 °C and reflux.


Said reaction is followed by the protection of the piperazine ring in a suitable solvent e.g. methylene chloride or chloroform with an appropriate protecting reagent e.g. di-*tert*-butyl dicarbonate with a suitable base e.g. triethylamine or K₂CO₃ and at a temperature between -20 °C and +60 °C.

(xii) Halogenation of the compound of formula **XV**, where R₁ is hydrogen, C₁-C₆-alkyl or C₃-C₆-cycloalkyl,



to obtain a compound of formula **XVII** may be performed by aromatic electrophilic substitution using a suitable halogenation agent such as Br₂, Cl₂, I₂, ICl, or SO₂Cl₂. The reaction may be carried out using the salt or the base of the compound **XV** in an appropriate solvent e.g. acetic acid, HCl/ethanol or water with or without a suitable base e.g. alkali metal acetate such as sodium acetate and at a reaction temperature between -20 °C and room temperature.


(xiii) Conversion of a compound of formula **XVII**, where R₁ is hydrogen, to a compound of formula **XVIII**,



where R_c is a suitable protecting group, may be carried out by the protection of the piperazine ring in a suitable solvent e.g. methylene chloride or chloroform with an

appropriate protecting reagent e.g. di-*tert*-butyl dicarbonate with a suitable base e.g. triethylamine or K_2CO_3 and at a temperature between $-20\text{ }^\circ\text{C}$ and $+60\text{ }^\circ\text{C}$.


(xiv) Halogenation of the compound of formula **XIX**, where R_2 is C_1-C_6 alkoxy (when X is O described in: Thorberg, S-O et al. Acta Pharm. Suec. 1987 24, 169-182; when X is CH_2 commercially available) either as racemate or as an enantiomer



(XIX)

(XX)


to obtain a compound of formula **XX** may be performed by aromatic electrophilic substitution using a suitable halogenation agent such as Br_2 , Cl_2 , I_2 , ICl , or SO_2Cl_2 . The reaction may be carried out using the salt or the base of the compound **XIX** in an appropriate solvent e.g. acetic acid, HCl /ethanol or water with or without a suitable base e.g. alkali metal acetate such as sodium acetate and at a reaction temperature between $-20\text{ }^\circ\text{C}$ and room temperature.




(XXI)

(xv) Benzylation of the compound of the formula **XX**, either as a racemate or as an enantiomer, to obtain a compound of the formula **XXI** by reaction with a suitable benzylation agent e.g. benzyl halide such as benzyl bromide or benzyl chloride or an

activated alcohol e.g. benzylmesylate or -tosylate. The reaction may be carried out using the salt or the base of compound of formula **XX** in a suitable solvent e.g. *N,N*-dimethylformamide, acetone or acetonitrile with a suitable base such as triethylamine, NaOH, NaHCO₃ or K₂CO₃ at a temperature within the range of +20 °C to +150 °C. The presence of a suitable catalyst e.g. alkali metal halide such as potassium iodide or sodium iodide may increase the speed of the reaction.




(xvi) Conversion of the compound of formula **XXI** to a compound of formula **XXII**, where R₁ is hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl and R₂ is C₁-C₆ alkoxy, may be carried out by the reaction with a compound of formula **XXIII**, where R₁ is hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl.



The process may be carried out in a suitable solvent e.g. an aprotic solvent such as benzene, toluene, dioxane, tetrahydrofuran or *N,N*-dimethylformamide with a suitable base such as sodium *tert*-butoxide or lithium bis(trimethylsilyl)amide in the presence of a suitable palladium catalyst such as PdZ₂, L'₂Pd(0) or L'₂PdZ₂ where Z stands for a


halogen such as chlorine or bromine and L' stands for a suitable ligand such as triphenylphosphine, tri-*o*-tolylphosphine, trifurylphosphine, triphenylarsine or dibenzylidenacetone and with or without an addition of a ligand L'' such as triphenylphosphine, tri-*o*-tolylphosphine, trifurylphosphine, 2,2'-bis(diphenylphosphino)-1,1'-binaphthalene (either as a racemate or as an enantiomer) or triphenylarsine and the reaction may occur at a temperature between +20 °C and +150 °C.

(xvii) Conversion of the compound of formula **XXII** to a compound of formula **XXIV**



where R₁ is hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl and R₂ is C₁-C₆ alkoxy may be carried out by hydrogenation using a catalyst containing palladium, platinum, rhodium or nickel in a suitable solvent e.g. acetic acid or ethanol at a reaction temperature between +20 °C and +120 °C.


(xviii) Conversion of compound of formula **XXIV**, where R₁ is hydrogen, to a compound of formula **XXV**,



(XXIV)

(XXV)

where R_c is a suitable protecting group, may be carried out by the protection of the
 5 piperazine ring in a suitable solvent e.g. methylene chloride or chloroform with a
 appropriate protecting reagent e.g. di-*tert*-butyl dicarbonate with a suitable base e.g.
 triethylamine or K_2CO_3 and at a temperature between $-20\text{ }^\circ\text{C}$ and $+60\text{ }^\circ\text{C}$.



(XV)


(XXVI)

(xix) Conversion of a compound of formula **XV**, where R_1 is C_1 - C_6 alkyl or C_3 - C_6
 10 cykloalkyl, to a compound of formula **XXVI**, where Y is $NHCO$ and R_3 is as defined in
 general formula **I** above, may be carried out by acylation with an appropriate benzoic acid
 of formula **XXVII** activated as an acid chloride in a suitable solvent such as methylene
 15 chloride or chloroform with a suitable base e.g. trialkylamine such as triethylamine or by

using a benzoic acid of formula **XXVII** with an activating reagent e.g. *N,N'*-carbonyldiimidazole, *N,N'*-dicyclohexylcarbodiimide or diphenylphosphinic chloride with a suitable base such as *N*-methylmorpholine in a suitable solvent such as *N,N*-dimethylformamide or tetrahydrofuran and the reaction may be conducted at a temperature
5 between +20 °C and +150 °C.

2. In the case where Y is CONH and X is CH₂ or O.


(i) Nitration of a compound of formula **XXVIII**, where R₂ is C₁-C₆ alkoxy, either as a racemate or as an enantiomer, to obtain a compound of formula **XXIX**,



where R_d is C₁-C₆ alkyl, may be carried out by aromatic electrophilic substitution using a
20 suitable nitration reagent such as nitric acid or nitric acid and sulphuric acid in a suitable solvent e.g. acetic acid, acetic anhydride or water at a reaction temperature between -20 °C and room temperature.


(ii) Hydrolysis of a compound of formula **XXIX** may be carried out under acidic
25 conditions using acids such as H₂SO₄, HCl, HBr, in a suitable solvent such as H₂O, ethanol, methanol, acetic acid or mixtures thereof and the reaction may occur at a temperature between +20 °C and reflux or under basic conditions using bases such as NaOH or KOH in a suitable solvent such as H₂O, ethanol, methanol or mixtures thereof

and the reaction may occur at a temperature between +20 °C and reflux, resulting in a compound of formula **XXX**.



(iii) Conversion of a compound of formula **XXX**, where R_2 is C_1 - C_6 alkoxy, to a compound of formula **XXXI**, where Y is CONH and R_2 is C_1 - C_6 alkoxy may be carried out by activation of the acid function of a compound of formula **XXX** as an acid halide such as an acid chloride with a suitable base such as a trialkylamine e.g. triethylamine or by using an activating reagent such as *N,N'*-carbonyldiimidazole, *N,N'*-dicyclohexylcarbodiimide or diphenylphosphinic chloride with a suitable base such as *N*-methylmorpholine in a suitable solvent e.g. methylene chloride, chloroform, toluene, *N,N*-dimethylformamide, dioxane or tetrahydrofuran followed by the addition of an appropriate aniline **XXXII**, where R_3 is as defined in formula **I** above. The reaction may occur between 0 °C and +120 °C.


(iv) Conversion of the compound of formula **XXXI** to a compound of formula **XXXIII**, where Y is CONH and R_3 is as defined in general formula **I** may be carried out by



(XXXIII)

5 hydrogenation using a catalyst containing palladium, platina or nickel in a suitable solvent such as ethanol, methanol or acetic acid at a reaction temperature between +20 °C and +120 °C; or reduction with sodium dithionite in a suitable solvent.

3. Conversion of a compound of formula XXXIV to a compound of formula XXXV



may be carried out by

15 a) hydrolysis of the nitrile in compound of formula XXXIV in a suitable solvent such as aqueous methanol or aqueous ethanol in the presence of a suitable base such as NaOH or KOH at a reaction temperature between room temperature and reflux, followed by


20 b) hydrolysis of the above formed amide and the ketal under acidic conditions in a suitable solvent such as aqueous methanol, aqueous ethanol or water in the presence of a suitable acid such as HCl or HBr at a reaction temperature between room temperature and reflux.

25 Methods of Preparation of End Products

Another object of the invention is a process A(i), A(ii), B or C for the preparation of the compound of general formula I by

A(i)

acylation, in the case where R₁ is C₁-C₆ alkyl or C₃-C₆ cycloalkyl, Y is NHCO and X, R₂ and R₃ are as defined in general formula I above, of a compound of formula A,




with an activated benzoic acid of formula **XXVII** or by using a benzoic acid of formula **XXVII** with an activating reagent.

Thus, the acylation according to the process A(i) may be carried out with an appropriate benzoic acid of formula XXVII, where R₃ is as defined in formula I above, activated as an acid chloride in a suitable solvent such as methylene chloride or chloroform with a suitable base e.g. trialkylamine such as triethylamine at a temperature between -20 °C and reflux temperature or by using an benzoic acid of formula XXVII, where R₃ is as defined in formula I above, with an activating reagent e.g. *N,N'*-carbonyldiimidazole, *N,N'*-dicyclohexylcarbodiimide or diphenylphosphinic chloride with a suitable base such as *N*-methylmorpholine in a suitable solvent such as *N,N*-dimethylformamide or tetrahydrofuran and the reaction may be conducted at a temperature between +20 °C and +150 °C.

A(ii)


acylation, in the case where R_1 is hydrogen, Y is NHCO, R_c is a protecting group and X, R_2 and R_3 are as defined in general formula I above, of a compound of formula B



Thus, the acylation according to the process A(ii) may be carried out with an appropriate benzoic acid of formula XXVII, where R_3 is as defined in formula I above, activated as an acid chloride in a suitable solvent such as methylene chloride or chloroform with a suitable base e.g. trialkylamine such as triethylamine at a temperature between $-20\text{ }^{\circ}\text{C}$ and reflux temperature or by using an benzoic acid of formula XXVII, where R_3 is as defined in formula I above, with an activating reagent e.g. *N,N'*-carbonyldiimidazole, *N,N'*-dicyclohexylcarbodiimide or diphenylphosphinic chloride with a suitable base such as *N*-methylmorpholine in a suitable solvent such as *N,N*-dimethylformamide or tetrahydrofuran and the reaction may be conducted at a temperature between $+20\text{ }^{\circ}\text{C}$ and $+150\text{ }^{\circ}\text{C}$ followed by removal of the protecting group R_c by hydrolysis in a suitable solvent such as methylene chloride or chloroform with a suitable acid such as trifluoroacetic acid at a temperature between $+20\text{ }^{\circ}\text{C}$ and $+60\text{ }^{\circ}\text{C}$.

B

reacting, in the case where Y is CONH, X, R₁, R₂ and R₃ are as defined in general formula I above,




with a compound of formula XI wherein L is a leaving group.

Thus, the reaction according to the process B may be carried out with a compound of formula XI wherein R₁ is as defined in general formula I above and L is a leaving group, e.g. a halogen such as chlorine or bromine or an alkane- or arenesulfonyloxy group such as p-toluene-sulfonyloxy group. The process may be carried out in a suitable solvent such as ethanol, butanol, *N,N*-dimethylformamide, acetonitrile or a mixture of water and acetonitrile with or without a suitable base e.g. K₂CO₃, NaHCO₃ or KOH and the reaction may occur between +20 °C and +150 °C.

C

reacting, in the case where Y is NHCO, R₂ is halogen and X, R₁ and R₃ are as defined in general formula I above, a compound of formula D



5 with a suitable halogenation agent such as Br₂, Cl₂, I₂, ICl, or SO₂Cl₂.

Thus, the reaction according to the process C may be carried out by aromatic electrophilic substitution using a suitable halogenation agent such as Br₂, Cl₂, I₂, ICl, or SO₂Cl₂. The reaction may be carried out using the salt or the base of the compound D in an appropriate solvent e.g. acetic acid, HCl/ethanol or water with or without a suitable base e.g. alkali metal acetate such as sodium acetate and at a reaction temperature between -20 °C and room temperature.

Working examples

15

Preparation of Intermediates and Starting Materials for the 5-HT_{1B}-antagonists

Preparation 1

(R)-2-N,N-Dibenzylamino-8-methoxy-1,2,3,4-tetrahydronaphthalene

20 To a solution of (R)-8-methoxy-2-amino-1,2,3,4-tetrahydronaphthalene hydrochloride (24 g, 0.11 mol) in acetonitrile (600 mL) were added potassium carbonate (53 g, 0.39 mol), potassium iodide (catalytic amount) and benzyl bromide (34 mL, 0.28 mol). The reaction mixture was stirred at reflux for a period of 35 h. After the precipitate was filtered off and the acetonitrile removed *in vacuo*, the residue was partitioned between diethyl ether and water. The organic phase was separated, dried (Na₂SO₄) and evaporated *in vacuo* to give a

25

crude product which was purified on a silica gel column using hexane/ethyl acetate, (3:1) as the eluent. Yield: 36 g (91%) of the title compound as a white solid: mp 105-107 °C; $[\alpha]^{21}_D +124^\circ$ (c 1.0, chloroform); EIMS (70 eV) m/z (relative intensity) 357 (100, M^+).

5 Preparation 2

(R)-7-N,N-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthol

(R)-2-N,N-Dibenzylamino-8-methoxy-1,2,3,4-tetrahydronaphthalene (43 g, 0.12 mol) was dissolved in diethyl ether (800 mL) and an excess of an ethereal HCl solution was added dropwise. The precipitate was filtered and dried *in vacuo* to give a white solid. This crude
10 product (42 g, 0.11 mol) was dissolved in anhydrous methylene chloride (1 L) and cooled to -60 °C. To the solution was boron tribromide (16 mL, 0.15 mol), dissolved in anhydrous methylene chloride (100 mL), added dropwise. The reaction temperature was allowed to reach -5 °C and was kept there overnight. To the ice-cooled solution was a 2 M aqueous ammonium hydroxide solution added dropwise and the mixture was extracted, twice, with
15 methylene chloride. The combined organic phases were dried (Na_2SO_4), filtered and the solvent removed *in vacuo* to give a crude residue. Chromatography on silica (eluent: methylene chloride) gave 34 g (93% yield) of the title compound as a viscous clear oil: $[\alpha]^{21}_D +118^\circ$ (c 1.5, chloroform); EIMS (70eV) m/z (relative intensity) 343 (53, M^+).

20 Preparation 3

(R)-2-(7-N,N-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthyloxy)-2-methylpropanamide

(R)-2-N,N-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthol (10 g, 29 mmol) was stirred in anhydrous dioxane (150 mL) with sodium hydride (80% in oil, 0.96 g, 32 mmol) for 1 h. 2-
25 Bromo-2-methylpropanamide (4.8 g, 29 mmol; described in: Coutts, I. G. C.; Southcott, M.R. *J. Chem. Soc. Perkin Trans. 1* 1990, 767-770) was added and the reaction mixture was heated at 100 °C for 2.5 h. After cooling, the precipitated sodium bromide was filtered off, the filtrate evaporated *in vacuo* and the residue was partitioned between water and methylene chloride. The organic phase was separated, dried (Na_2SO_4), filtered and
30 evaporated to give a crude product which was purified on a silica gel column using methylene chloride as the eluent. Yield: 9.6 g (76%) of the title compound as white

crystals: mp 125-126 °C; $[\alpha]^{21}_D +98^\circ$ (c 1.1, chloroform); EIMS (70eV) m/z (relative intensity) 428 (13, M^+).

Preparation 4

5 **(R)-N-(7-N,N-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthyl)-2-hydroxy-2-methylpropanamide**

To a solution of (R)-2-(7-N,N-dibenzylamino-5,6,7,8-tetrahydro-1-naphthyloxy)-2-methylpropanamide (9.1 g, 21 mmol) in anhydrous 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone (10 mL) and dry N,N-dimethylformamide (100 mL) was added sodium hydride
10 (80% in oil, 1.4 g, 47 mmol) and the reaction was heated at 130 °C for 8 h. The solution was poured into a mixture of ice and water and extracted three times with ethyl acetate. The combined organic phases were dried (Na_2SO_4), filtered and evaporated *in vacuo*. Chromatography on silica (eluent: chloroform/ethanol saturated with NH_3 ; 100:0.5) gave 7.6 g (84% yield) as white crystals: mp 134-135 °C; $[\alpha]^{21}_D +130^\circ$ (c 1.1, chloroform);
15 EIMS (70eV) m/z (relative intensity) 428 (1, M^+).

Preparation 5

(R)-2-N,N-Dibenzylamino-8-amino-1,2,3,4-tetrahydronaphthalene

(R)-N-(7-N,N-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthyl)-2-hydroxy-2-
20 methylpropionamide (7.4 g, 17 mmol) was dissolved in a mixture of ethanol (200 mL) and a 20% HCl aqueous solution (300 mL) and heated to reflux for 8 h. The ethanol was evaporated *in vacuo* and the remaining solution was washed twice with diethyl ether and cooled on ice-bath. After alkalization with a 45% aqueous solution of sodium hydroxide the mixture was extracted with methylene chloride. The combined organic phases were
25 dried (Na_2SO_4), filtered and evaporated *in vacuo*. Purification on a silica gel column using chloroform as the eluent gave 3.8 g (76% yield) of the title compound as a light-brown oil: $[\alpha]^{21}_D +124^\circ$ (c 0.9, chloroform); EIMS (70eV) m/z (relative intensity) 342 (92, M^+).

Preparation 6**(R)-1-(7-*N,N*-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthyl)-4-*N*-methylpiperazine-2,6-dione**

1,1'-Carbonyldiimidazole (6.0 g, 37 mmol) was added to a stirred suspension of methyliminodiacetic acid (2.7 g, 18 mmol) in anhydrous tetrahydrofuran (250 mL). The reaction mixture was heated at reflux for 1.5 h. (R)-2-*N,N*-Dibenzylamino-8-amino-1,2,3,4-tetrahydronaphthalene (5.7 g, 17 mmol) was then added and stirring at reflux was continued for 17 h. An additional amount of 1,1'-carbonyldiimidazole (2.9 g, 18 mmol) was added and heating at reflux was continued for another 17 h. The solvent was evaporated *in vacuo* and the crude product was purified on a silica gel column using chloroform/ethanol saturated with NH₃ (100:0.5) as the eluent. Yield: 6.6 g (87%) of the title compound as an oil: $[\alpha]^{21}_{\text{D}} +90^{\circ}$ (*c* 0.52, chloroform); EIMS (70eV) *m/z* (relative intensity) 453 (8, M⁺).

Preparation 7

(R)-2-*N,N*-Dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (R)-1-(7-*N,N*-Dibenzylamino-5,6,7,8-tetrahydro-1-naphthyl)-4-methylpiperazine-2,6-dione (1.4 g, 3.1 mmol) was added to a suspension of lithium aluminium hydride (0.57 g, 15 mmol) in anhydrous diethyl ether (70 mL). The reaction mixture was heated at reflux for 7 h. The reaction was quenched by the addition of water (0.60 mL), 15% aqueous sodium hydroxide (0.60 mL) and again water (1.8 mL). The mixture was filtered, dried (Na₂SO₄) and evaporated *in vacuo*. Purification on a silica gel column using chloroform/ethanol saturated with NH₃ (100:2) as the eluent gave 1.0 g (79% yield) of the title compound as a viscous oil: $[\alpha]^{21}_{\text{D}} +53^{\circ}$ (*c* 0.5, chloroform); EIMS (70 eV) *m/z* (relative intensity) 425 (2, M⁺).

Preparation 8**(R)-5-Bromo-2-*N,N*-dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene.**

To a solution of (R)-2-*N,N*-dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (2.8 g, 6.5 mmol) and sodium acetate (6.8 g, 83 mmol) in acetic acid

(100 mL) bromine (370 μ L, 7.2 mmol) was added in one portion and the reaction was stirred for 5 min. The solvent was evaporated *in vacuo* and the remaining solid was partitioned between water and methylene chloride and cooled on ice-bath. The water phase was alkalized with 2 M aqueous solution of sodium hydroxide and the phases were
5 separated. The organic phase was dried (Na_2SO_4), filtered and evaporated *in vacuo* to give a crude product which was purified on a silica gel column using chloroform/ethanol saturated with NH_3 (100:2) as the eluent. Yield: 2 g (61%) of a viscous brown oil: EIMS (70 eV) m/z (relative intensity) 503 and 505 (0.6, M^+)

10 Preparation 9

(R)-2-N,N-Dibenzylamino-8-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalene

(R)-2-N,N-Dibenzylamino-8-amino-1,2,3,4-tetrahydronaphthalene (9.8 g, 39 mmol) and bis-(2-chloroethyl)amine hydrochloride (5.5 g, 32 mmol) was dissolved in n-butanol (80 mL). The reaction mixture was stirred at 100 $^\circ\text{C}$ and after 65 h the mixture was filtered and
15 the solvent evaporated *in vacuo*. Purification on a silica gel column using chloroform/methanol/ concentrated ammonium hydroxide (95:5:0.5) as the eluent gave 6.0 g (51% yield) of the title compound as a viscous oil: $[\alpha]^{21}_{\text{D}} +72^\circ$ (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 411 (2, M^+).

20 Preparation 10

(R)-2-Amino-8-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalene

To a solution of (R)-2-N,N-dibenzylamino-8-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (5.5 g, 13 mmol) in methanol (400 mL) were added ammonium formate (20 g, 0.32 mol) and palladium (10%) on activated carbon (1.9 g). The mixture
25 was refluxed for 1 h and the palladium was then filtered off. The solvent was evaporated *in vacuo* and the residue was partitioned between methylene chloride and a 2 M ammonium hydroxide solution. The organic phase was separated, dried (Na_2SO_4), filtered and evaporated *in vacuo* to give a crude product which was purified on a silica gel column using chloroform/ethanol/concentrated ammonium hydroxide (80:20:2.5) as the eluent.

Yield: 2.4 g (76%) of the title compound as an oil: $[\alpha]^{21}_D +9.9^\circ$ (c 1.0, chloroform);
EIMS (70 eV) m/z (relative intensity) 231 (24, M^+).

Preparation 11

5 **(*R*)-2-Amino-5-bromo-8-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalene**

The title compound was prepared from (*R*)-2-amino-8-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalene following the general method of preparation 8. Purification on a silica gel column using methylene chloride/ethanol/concentrated ammonium hydroxide (80:20:2) as the eluent gave 0.8 g (67% yield) of a viscous light brown oil: $[\alpha]^{21}_D -6.2^\circ$
10 ($c=1$, chloroform); EIMS (70 eV) m/z (relative intensity) 309 and 311 (3.5, M^+)

Preparation 12

***tert*-Butyl (*R*)-4-(7-Amino-4-bromo-5,6,7,8-tetrahydro-1-naphthyl)piperazin-1-carboxylate**

15 To an ice-cooled solution of (*R*)-2-amino-5-bromo-8-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (0.8 g, 2.6 mmol) and triethylamine (0.53 mL, 3.9 mmol) in methylene chloride (50 mL) was added di-*tert*-butyl dicarbonate (0.56 g, 2.6 mmol) dissolved in methylene chloride (10 mL). After the addition, the reaction was allowed to stir at ambient temperature for 1 h. Water (10 mL) was added and the mixture was cooled
20 on an ice-bath. The water phase was alkalized with a 2 M aqueous solution of sodium hydroxide and the phases were separated. The organic phase was dried (Na_2SO_4), filtered and evaporated *in vacuo* to give a crude product which was purified on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (95:5:0.5) as the eluent.
Yield: 0.41 g (38%) of a viscous colorless oil: $[\alpha]^{21}_D +13^\circ$ ($c=1$, chloroform); EIMS (70
25 eV) m/z (relative intensity) 409 and 411 (75, M^+)

Preparation 13

(*R*)-*N*-[5-Bromo-8-(4-*tert*-butyloxycarbonylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

30 4-Morpholinobenzoic acid (0.50 g, 2.4 mmol; described in: Degutis, J.;

Rasteikiene, L.; Degutiene, A. *Zh. Org. Khim.* **1978**, *14*(10), 2060-2064) was dissolved in thionyl chloride (10 mL). After 2 min, the thionyl chloride was evaporated *in vacuo* and the residue was treated with toluene and again the solvent was evaporated *in vacuo*. Crude acid chloride (81 mg, 0.36 mmol) was dissolved in methylene chloride (10 mL) and added dropwise to a solution of *tert*-butyl (*R*)-4-(7-amino-4-bromo-5,6,7,8-tetrahydro-1-naphthyl)piperazin-1-carboxylate (140 mg, 0.34 mmol) and triethylamine (71 μ L, 0.51 mmol) in methylene chloride (10 mL). After the addition, the reaction was stirred at ambient temperature for 15 min and was then washed with a diluted aqueous solution of sodium hydrogen carbonate and the phases were separated. The organic phase was dried (Na_2SO_4), filtered and evaporated *in vacuo* and the residue was purified on a silica gel column using chloroform/ethanol saturated with NH_3 (100:2) as the eluent. Yield: 160 mg (79%) of a viscous colorless oil: $[\alpha]_D^{21} -11^\circ$ ($c=1$, chloroform); TSPMS m/z (relative intensity) 599 and 601 (35, $\text{M}^+ + 1$).

Preparation 14

(*R*)-2-Amino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene

To a solution of (*R*)-2-*N,N*-dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (4.0 g, 9.4 mmol) in methanol (250 mL) were added ammonium formate (14 g, 56 mmol) and palladium (10%) on activated carbon (1.4 g). The mixture was refluxed for 3 h and the palladium was then filtered off. The solvent was evaporated *in vacuo* and the residue was partitioned between methylene chloride and a 2 M ammonium hydroxide solution. The organic phase was separated, dried (Na_2SO_4), filtered and evaporated *in vacuo* to give a crude product which was purified on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (90:9:0.5) as the eluent. Yield: 1.9 g (83%) as an oil: $[\alpha]_D^{21} -2.7^\circ$ (c 1.0, chloroform); EIMS (70 eV) m/z (relative intensity) 245 (5, M^+).

Preparation 15**(R)-2-Amino-5-bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene**

The title compound was prepared from (R)-2-amino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene following the general method of Preparation 8. Purification on a silica gel column using chloroform/ethanol/concentrated ammonium hydroxide (80:20:2) as the eluent gave 630 mg (89% yield) of a viscous colorless oil: EIMS (70 eV) *m/z* (relative intensity) 323 and 325 (20, M⁺)

Preparation 16**(R)-2-Amino-8-bromo-5-methoxy-1,2,3,4-tetrahydronaphthalene Hydrochloride**

(R)-2-Amino-5-methoxy-1,2,3,4-tetrahydronaphthalene hydrochloride (5.0 g, 23 mmol) was dissolved in acetic acid (300 mL) under nitrogen atmosphere. Sodium acetate (5.5 g, 70 mmol) was added and bromine (3.5 g, 23 mmol) was then added in one portion. The mixture was stirred for 5 minutes at room temperature. The solvent was removed *in vacuo* to give a solid residue which was partitioned between ethyl acetate and NaOH (2 M). The layers were separated and the aqueous phase was extracted twice with ethyl acetate. The organic layers were combined and dried (Na₂SO₄). The solvent was removed *in vacuo* to give a brown oily residue. The HCl salt was precipitated from diethyl ether/methylene chloride by the addition of HCl in diethyl ether (3 M): yield 7.7 g (94%). Recrystallization from methanol gave the title compound as needle crystals: mp 264-265 °C; [α]_D²¹ +54° (c 1, MeOH); EIMS (70eV) *m/z* (relative intensity) 257 (30, M⁺, ⁸¹Br), 255 (31, M⁺, ⁷⁹Br).

Preparation 17**(R)-8-Bromo-2-N,N-dibenzylamino-5-methoxy-1,2,3,4-tetrahydronaphthalene**

(R)-2-Amino-8-bromo-5-methoxy-1,2,3,4-tetrahydronaphthalene hydrochloride (4.5 g, 17.5 mmol), benzyl bromide (6.6 g, 38 mmol), potassium carbonate (9.7 g, 70 mmol) and potassium iodide (100 mg, catalytic amount) were mixed with acetonitrile (250 mL) under nitrogen atmosphere and refluxed for 18 h. The solvent was removed *in vacuo* and the residue was partitioned between ethyl acetate and ammonia (2 M). The layers were separated and the organic layer was dried (MgSO₄). The solvent was removed *in vacuo* to

give a residue which was purified by flash chromatography on silica gel using hexane/methylene chloride 8:2 as the eluent. The title compound was obtained as an oil. Yield 7.5 g (98%): $[\alpha]^{21}_{\text{D}} + 87^\circ$ (c 1, MeOH); EIMS (70eV) m/z (relative intensity) 437 (12, M^+ , ^{81}Br), 435 (13, M^+ , ^{79}Br).

5

Preparation 18

(*R*)-2-*N,N*-Dibenzylamino-5-methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene

To a solution of (*R*)-8-bromo-2-*N,N*-dibenzylamino-5-methoxy-1,2,3,4-tetrahydronaphthalene (19 g, 44 mmol) in dry toluene (500 mL) under an argon atmosphere was added *N*-methylpiperazine (5.9 mL, 53 mmol), tris(dibenzylideneacetone)dipalladium(0) (0.41 g, 0.44 mmol), (*R*)-BINAP (0.82 g, 1.3 mmol) and sodium *tert*-butoxide (0.40 mg, 4.2 mmol). The dark solution was stirred at 85 °C for 23 h and was then cooled, filtered and evaporated *in vacuo*. Purification on a silica gel column using chloroform/ethanol saturated with NH_3 (100:2) as the eluent gave 19 g (97% yield) of a viscous colorless oil: $[\alpha]^{21}_{\text{D}} + 72^\circ$ ($c=1$, chloroform); EIMS (70eV) m/z (relative intensity) 455 (15, M^+).

15

Preparation 19

(*R*)-2-Amino-5-methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene

The title compound was prepared from (*R*)-2-*N,N*-dibenzylamino-5-methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene following the general method of Preparation 10. Yield: 5.3 g (82%) of a viscous colorless oil: $[\alpha]^{21}_{\text{D}} + 20^\circ$ ($c=1.1$, chloroform); EIMS (70eV) m/z (relative intensity) 275 (53, M^+).

25

Preparation 20

Methyl 5-Methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxylate

Methyl 5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxylate (1.1 g, 5 mmol; described in: Johnson, D.W.; Mander, L.N. *Aust.J.Chem.* 1974, 8, 1277-1286) dissolved in acetic anhydride (20 mL), was treated with 70% nitric acid (0.4 mL) at 0 °C for 1 h and the

30

mixture was poured into ice-water and diethyl ether. The organic phase was separated, evaporated *in vacuo* and the residue triturated with diisopropyl ether to yield 0.27 g (20%) of the title compound as crystals: mp 100-104 °C; EIMS (70 eV) *m/z* (relative intensity) 265 (35, M⁺).

5

Preparation 21

5-Methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxylic Acid

A mixture of methyl 5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxylate (1.9 g, 7.1 mmol) in methanol (20 mL) and 2 M NaOH (10 mL) was refluxed for 1.5 h and the solvent was evaporated *in vacuo*. The residue was taken up in ethyl acetate and acidified. The organic phase was separated and dried and evaporated *in vacuo* to afford 1.7 g (95% yield) of crystals: mp (after recrystallization in diisopropyl ether/ethanol) 189-190 °C; EIMS (70eV) *m/z* (relative intensity) 251 (30, M⁺).

Preparation 22

***N*-(4-Morpholinophenyl)-5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxamide**

A mixture of 5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxylic acid (1.3 g, 5 mmol), toluene (20 mL) and thionyl chloride (1.8 mL, 25 mmol) was heated at 80 °C for 1 h. The solvents were removed *in vacuo* and the residue, dissolved in methylene chloride (10 mL), was added to a solution of 4-morpholinoaniline (890 mg, 5 mmol) and triethylamine (1.0 g, 10 mmol) in methylene chloride (20 mL) at 0 °C. The mixture was stirred at 20 °C for 2 h, water was added and the precipitate was filtered to yield 1.9 g (90%) of the title product as crystals: mp 251-253 °C; EIMS (70 eV) *m/z* (relative intensity) 411 (100, M⁺).

Preparation 23

***N*-(4-Morpholinophenyl)-8-amino-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide**

A solution of *N*-(4-morpholinophenyl)-5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxamide (2.05 g, 5 mmol) and sodium dithionite (3.5 g, 20 mmol) in *N,N*-

dimethylformamide (20 mL) and water (2 mL) was heated at 90 °C for 7 h. After cooling, the reaction mixture was partitioned between water and ethyl acetate, the phases were separated and the organic phase was washed, twice, with water and evaporated *in vacuo*. The residue was triturated with diisopropyl ether/ethyl acetate affording 1.4 g (72% yield) of the title product as crystals: mp 219-222 °C; EIMS (70eV) *m/z* (relative intensity) 381 (70, M⁺).

Preparation 24

***N*-(4-Morpholinocarbonylphenyl)-5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxamide**

A mixture of 5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxylic acid (1.0 g, 4 mmol), toluene (20 mL), *N,N*-dimethylformamide (10 drops) and thionyl chloride (1.5 mL, 20 mmol) was heated at 60 °C for 1 h. The solvents were removed *in vacuo* and the residue, dissolved in methylene chloride (20 mL), was added to a solution of 4-aminobenzoylmorpholine (820 mg, 4 mmol, described in: Devlin J.P. *J.Chem.Soc. Perkin Trans I*, 1975, 830-841) and triethylamine (800 mg, 8 mmol) in methylene chloride (30 mL) at 5 °C. After stirring at 20 °C for 2 h, water was added and the organic phase was separated, dried and the solvent removed *in vacuo*. The oily residue was crystallized from diisopropyl ether/ethyl acetate affording 1.2 g (73% yield) of the title compound as crystals: mp 186-189 °C; EIMS (70eV) *m/z* (relative intensity) 439 (20, M⁺).

Preparation 25

***N*-(Morpholinocarbonylphenyl)-8-amino-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide**

A solution of *N*-(4-morpholinocarbonylphenyl)-5-methoxy-8-nitro-1,2,3,4-tetrahydronaphthalene-2-carboxamide (1.3 g, 2.8 mmol) and sodium dithionite (2.0 g, 11 mmol) in *N,N*-dimethylformamide (20 mL) and water (2.5 mL) was heated at 85 °C for 3 h. After cooling, the reaction mixture was partitioned between water and ethyl acetate, the phases were separated and the organic phase was washed, twice, with water and evaporated *in vacuo*. The organic phase was dried and evaporated. The residue was treated with

diisopropyl ether affording 310 mg (30% yield) of the title product as crystals: EIMS (70eV) m/z (relative intensity) 409 (100, M^+).

Preparation 26

5 **(*R*)-2-*N,N*-Dibenzylamino-5-(1-hydroxyethyl)-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene**

(*R*)-5-Bromo-2-*N,N*-dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (1.4 g, 2.8 mmol) was dissolved in freshly distilled tetrahydrofuran (100 mL), flushed with argon and cooled to -78 °C. To the solution was added *tert*-butyl lithium (2.6 mL, 1.4 M in pentane, 3.7 mmol) and the reddish solution was stirred at ambient
10 temperature for 10 min. Acetaldehyde (320 μ L, 5.7 mmol) was added and the reaction mixture was stirred at -78 °C for 10 min, at 0 °C for 2 h and at room temperature for 10 min. The reaction was quenched with water and the solvent was evaporated *in vacuo*. The residue was partitioned between diethyl ether (100 mL) and 2 M NH_3 (20 mL) and the
15 aqueous phase was extracted with diethyl ether (20 mL). The combined organic layers were washed with brine (20 mL) and dried ($MgSO_4$). The solvent was evaporated giving 2.0 g of a crude product. Purification by column chromatography on silica gel using chloroform/methanol/conc. NH_3 (95:5:0.5) as the eluent gave 910 mg (68% yield) of the title compound as a yellowish foam: ESI m/z (relative intensity) 470 (100, $M+1$).

20

Preparation 27

(*R*)-2-Amino-5-ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene

(*R*)-2-*N,N*-Dibenzylamino-5-(1-hydroxyethyl)-1,2,3,4-tetrahydronaphthalene (1.6 g, 3.4 mmol) was dissolved in acetic acid (80 mL) and stirred at 100 °C for 2 h. The solvent was
25 evaporated *in vacuo* and the residue was dissolved in methanol (150 mL). Palladium (10%) on charcoal (600 mg) was added and the solution was flushed with nitrogen. To the solution was added ammonium formate (1.7 g, 28 mmol) and the reaction mixture was stirred at 65 °C for 2 h. The catalyst was filtered off and the solvent was evaporated *in vacuo* giving 1.3 g of a crude product. The residue was partitioned between methylene
30 chloride (120 mL) and 2 M NH_3 (30 mL). The organic phase was washed with brine (20 mL) and dried ($MgSO_4$). The solvent was evaporated *in vacuo* giving 740 mg (79% yield)

of the title compound as a white semi-crystalline solid: EIMS (70 eV) m/z (relative intensity) 273 (24, M^+).

Preparation 28

5 **(*R*)-*N*-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-trifluoromethylbenzamide**

To an ice-cooled solution of (*R*)-2-amino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (110 mg, 0.44 mmol) and triethylamine (91 μ L, 0.66 mmol) in methylene chloride (20 mL) was 4-(trifluoromethyl)benzoyl chloride (96 mg, 0.46 mmol)
10 in methylene chloride (5 mL) added dropwise. After the addition the reaction was allowed to stir at ambient temperature for 15 min and was then washed with diluted aqueous sodium hydrogen carbonate. The phases were separated and the organic phase was dried (Na_2SO_4), filtered and evaporated *in vacuo* to give a crude product which was purified on a silica gel column using chloroform/ethanol saturated with NH_3 (100:2) as the eluent.
15 Yield: 150 mg (81%) of the title compound as white crystals: mp 203-204 °C; $[\alpha]^{21}_{\text{D}} -20^\circ$ (*c* 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 417 (10, M^+).

Preparation 29

20 **(*R*)-2-*N,N*-Dibenzylamino-5-hydroxymethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene**

(*R*)-5-Bromo-2-*N,N*-dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (800 mg, 1.6 mmol) was dissolved in freshly distilled tetrahydrofuran (80 mL), flushed with argon and cooled to -78 °C. To the solution was added *tert*-butyl lithium (1.5 mL, 1.4 M in pentane, 2.1 mmol) and the reaction mixture
25 was stirred at ambient temperature for 10 min. Methyl chloroformate (250 μ L, 3.2 mmol) was added and the reaction mixture was stirred at -78 °C for 50 min and at 0 °C for 1 h. The reaction was quenched with water and the solvent was evaporated *in vacuo*. The residue was partitioned between diethyl ether (90 mL) and 2 M NH_3 (15 mL). The organic layer was washed with brine (10 mL) and dried (MgSO_4). The solvent was evaporated *in vacuo* giving 770 mg of a crude product. Purification by column chromatography on silica
30 gel using chloroform/methanol/conc. NH_3 (250:5:0.5) as the eluent afforded 610 mg of

(*R*)-5-carboxymethyl-2-*N,N*-dibenzylamino-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (containing 13% of the corresponding 5-hydrogen analogue) as a yellowish oil: EIMS (70 eV) *m/z* (relative intensity) 483 (1, M^+). The methyl ester (610 mg, 1.1 mmol) was dissolved in freshly distilled tetrahydrofuran (35 mL) and lithium aluminum hydride (120 mg, 3.1 mmol) was added. The reaction mixture was stirred at 45 °C for 2 h followed by cooling to room temperature. The reaction was quenched with water (120 μ L), 15% NaOH (120 μ L) and water (240 μ L) followed by stirring the slurry at room temperature for 2.5 h. The precipitate was filtered off and the solvent was evaporated *in vacuo* giving 730 mg of a crude product. Purification by column chromatography on a silica gel column using chloroform/methanol/conc. NH_3 (95:5:0.5) as the eluent gave 360 mg (50% yield) of the title compound as a white foam: EIMS (70 eV) *m/z* (relative intensity) 455 (1, M^+); $[\alpha]^{21}_D +44^\circ$ (*c* 0.12, chloroform).

Preparation 30

(*R*)-2-Amino-5-methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (*R*)-2-*N,N*-Dibenzylamino-5-hydroxymethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (360 mg, 0.78 mmol) was dissolved in methanol (35 mL), palladium (10%) on charcoal (170 mg) was added and the solution was flushed with nitrogen. To the solution was added ammonium formate (390 mg, 6.2 mmol) and the reaction mixture was stirred at 65 °C for 13 h. The catalyst was filtered off and the solvent was evaporated *in vacuo* giving 220 mg of a residue. The crude hydroxymethyl compound was dissolved in acetic acid (25 mL), palladium (10%) on charcoal (60 mg) was added and the solution was flushed with hydrogen. The reaction mixture was hydrogenated at room temperature and at atmospheric pressure for 4 h. The catalyst was filtered off and more palladium (10%) on charcoal (160 mg) was added followed by hydrogenation at room temperature and at atmospheric pressure for 24 h. The catalyst was filtered off and the solvent was evaporated *in vacuo*. The residue was partitioned between diethyl ether (70 mL) and conc. NH_3 and the organic phase was washed with brine (5 mL). The organic layer was dried ($MgSO_4$) and the solvent was evaporated *in vacuo* to give 120 mg (61% yield) of the title compound as a white semi-crystalline solid: EIMS *m/z* (relative intensity) 259 (20, M^+); $[\alpha]^{21}_D -1^\circ$ (*c* 0.09, chloroform).

Preparation 31

(S)-3-*N,N*-Dibenzylamino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran hydrochloride.

(S)-3-Amino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran (45 g, 0.25 mol; described in WO 93/07135), K₂CO₃ (120 g, 0.87 mol) and benzylbromide (65 mL, 0.55 mol) were mixed in acetonitrile (1000 mL) under nitrogen. The reaction mixture was refluxed for 45 h. The mixture was filtered and the solvent was removed *in vacuo*, and the residue was partitioned between diethyl ether and saturated NaCl (aq). The layers were separated and the organic phase was dried (MgSO₄) and filtered followed by precipitation of the hydrochloric salt at room temperature. Yield: 99 gram (99%). An analytical sample was transferred to the base: $[\alpha]^{21}_{\text{D}} +116^{\circ}$ (*c* 1.0, chloroform). EIMS (70eV) *m/z* (relative intensity) 359 (28, M⁺).

Preparation 32

(S)-3-*N,N*-Dibenzylamino-5-hydroxy-3,4-dihydro-2*H*-1-benzopyran.

(S)-3-*N,N*-Dibenzylamino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran hydrochloride (67 g, 0.17 mol) was dissolved in methylene chloride (500 mL) under nitrogen, and the solution was cooled to -75 °C. Boron tribromide (32 mL, 0.34 mol) was added dropwise over 5 min. The temperature was then allowed to slowly reach +5 °C, and the reaction was stirred over night. The reaction mixture was carefully quenched with an 2 M aqueous solution of NH₃ under stirring. The layers were separated and the aqueous phase was extracted two times with methylene chloride. The organic layers were combined, washed with brine, dried (MgSO₄), filtered and the solvent was removed *in vacuo* to give a brownish oily residue which was purified by flash chromatography on a silica gel column using methylene chloride as the eluent. Yield: 50 g (86%) of the title compound: $[\alpha]^{21}_{\text{D}} +109^{\circ}$ (*c* 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 345 (5, M⁺).

Preparation 33

(S)-2-(3-*N,N*-Dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yloxy)-2-methylpropanamide.

(S)-3-*N,N*-Dibenzylamino-5-hydroxy-3,4-dihydro-2*H*-1-benzopyran (50 g, 0.14 mol) was dissolved in anhydrous 1,4-dioxane (450 mL) under nitrogen. A dispersion of sodium

hydride (60-65% in oil, 6.1 g, 0.15 mol) was added in portions. The mixture was stirred for 1 h at room temperature. 2-Bromo-2-methylpropanamide (24 g, 0.14 mol; Coutts, I. G. C.; Southcott, M. R. *J. Chem. Soc. Perkin Trans. 1* 1990, 767-771) was added to the dark greenish solution and was heated at reflux with stirring for 3 h. An additional amount of sodium hydride (60-65% in oil, 2.8 g, 70 mmol) and 2-bromo-2-methylpropanamide (4.6 g, 28 mmol) was added in portions and heating at 60 °C was continued for 17 h. After cooling, a small amount of methanol (10 mL) was added and the solution was filtered and the solvent was removed *in vacuo*. The residue was partitioned between ethyl acetate (500 mL) and a saturated NaHCO₃ solution (50 mL). The organic layer was dried (MgSO₄), and the solvent was removed *in vacuo* to give a brownish residue which was crystallized from ethyl acetate/hexane. Yield: 45 g (71%) of the title compound as a white solid: mp 133-134 °C; [α]_D²¹ +99° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 430 (9, M⁺).

Preparation 34

(*S*)-5-Amino-3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran.

To a solution of (*S*)-2-(3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yloxy)-2-methylpropanamide (46 g, 0.11 mol) in anhydrous *N,N*-dimethylformamide (450 mL) and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (45 mL) was added sodium hydride (60-65% in oil, 8.5 g, 0.21 mol) in portions under nitrogen. The reaction mixture was heated at 110 °C with stirring for 13 h. The mixture was then allowed to cool, and the solution was partitioned between ethyl acetate (400 mL) and a 2 M NH₃ solution (200 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (150 mL). The combined organic layers were dried (MgSO₄) and concentrated *in vacuo* to give a brownish oil. EIMS (70eV) *m/z* (relative intensity) 430 (3, M⁺). The obtained material (0.11 mol) was dissolved in ethanol (350 mL). A 6 M HCl solution (250 mL) was added, and the reaction mixture was heated at reflux for 16 h. After stirring, the mixture was allowed to cool to 35 °C, the ethanolic solvent was removed *in vacuo*, and ethyl acetate was added to the aqueous remains. The mixture was cooled on ice, and a solution of conc. NH₃ was slowly added with stirring. The layers were separated, and the aqueous layer was extracted with another portion of ethyl acetate. The combined organic layers

were dried (MgSO_4), and the solvent was removed *in vacuo* to give a brownish oil which was purified on a short column of silica gel (eluent: hexane/ethyl acetate; 8:2) affording 25 g (68% yield) of the desired compound as a light yellow oil. The product slowly crystallized upon standing in the refrigerator. An analytical sample was recrystallized from diethyl ether/petroleum ether: mp 101-103 °C; $[\alpha]^{21}_{\text{D}} +123^\circ$ (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 344 (17, M^+).

Preparation 35

(S)-1-(3-*N,N*-Dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yl)-4-methylpiperazine-2,6-dione.

To a dispersion of *N*-methyliminodiacetic acid (6.90 g, 46.9 mmol) in anhydrous tetrahydrofuran (575 mL) was added 1,1'-carbonyldiimidazole (15.2 g, 93.9 mmol), and the mixture was heated at reflux for 2 h under nitrogen. A solution of (*S*)-5-amino-3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran (15.0 g, 42.7 mmol) in tetrahydrofuran (120 mL) was added with stirring over 0.5 h. The reaction mixture was heated at reflux for 28 h, then allowed to cool, and the solvent was removed *in vacuo*. The residue was purified on a short column of silica gel (eluent: methylene chloride and ethyl acetate) affording 14.1 g (71% yield) of the title compound as a light yellow solid: mp sinters >60 °C; $[\alpha]^{21}_{\text{D}} +89^\circ$ (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 455 (8, M^+).

Preparation 36

(S)-3-*N,N*-Dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran.

To a stirred solution of (*S*)-1-(3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yl)-4-methylpiperazine-2,6-dione (25.4 g, 55.8 mmol) in anhydrous diethyl ether (800 mL) was added lithium aluminum hydride (9.30 g, 246 mmol) in portions. The reaction mixture was heated to reflux for 6.5 h under nitrogen and was stirred over night at room temperature. The mixture was cooled (ice-bath), and water (10 mL) was added followed by a 15% aqueous solution of NaOH (10 mL) and another portion of water (30 mL). The precipitate was filtered off and washed with several portions of warm tetrahydrofuran. The organic layers were combined, and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 95:5 + 0.5% conc. NH_3)

affording 13.6 g (57% yield) of the title compound as a light yellow oil: $[\alpha]_D^{25} +63^\circ$ (*c* 1.0, methanol); EIMS (70eV) *m/z* (relative intensity) 427 (5, M^+).

Preparation 37

(*S*)-3-Amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran.

To a solution of (*S*)-3-*N,N*-dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran (2.6 g, 6.2 mmol) in anhydrous methanol (100 mL) were added palladium (10%) on activated carbon (0.97 g) and ammonium formate (3.1 g, 49 mmol) under nitrogen. The reaction mixture was heated at 50 °C with stirring overnight. The solution was filtered through Celite[®], and the solvent was removed *in vacuo*. The residue was partitioned between a 2 M NH_3 solution (20 mL) and ethyl acetate (100 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic phases were dried (Na_2SO_4), and the solvent was removed *in vacuo* to give 1.4 g (89% yield) of the title compound as a pale yellow oil: $[\alpha]_D^{21} -15^\circ$ (*c* 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 247 (74, M^+).

Preparation 38

4-(4-Piperidon-1-yl)benzoic Acid.

A solution of 2 M NaOH (10 mL), 4-(8-aza-1,4-dioxaspiro[4,5]dec-8-yl)benzonitrile (820 mg, 3.36 mmol; described in: Taylor E. C.; Skotnicki J. S. *Synthesis* **1981**, 8, 606-608), and ethanol (7.5 mL) was heated at reflux for 3 h. The external heating was interrupted, and the reaction mixture was stirred overnight at ambient temperature. The ethanolic solvent was removed *in vacuo*, and the remains were acidified to pH 4 with a 2 M HCl solution followed by extraction with ethyl acetate (50 mL). The layers were separated, and pH was adjusted to pH 6 with a 2 M NaOH solution followed by another extraction with ethyl acetate (50 mL). The combined organic layers were concentrated *in vacuo*, and the solid residue was dissolved in a 6 M HCl solution (10 mL). The reaction mixture was heated at 75 °C for 2.5 h and then at 55 °C overnight. The temperature was raised to 75 °C for 2 h, and the reaction mixture was then allowed to cool. The pH was adjusted to pH 4, and the solution was extracted with ethyl acetate (50 mL). The layers were separated, and another extraction was made at pH 5. The combined organic layers were dried ($MgSO_4$), and the

solvent was removed *in vacuo*. The crude product was recrystallized from ethyl acetate affording 300 mg (41% yield) of the title compound as yellowish crystals: mp sinters >215 °C; EIMS (70eV) *m/z* (relative intensity) 219 (100, M⁺)

5 **Preparation 39**

(S)-3-*N,N*-Dibenzylamino-8-iodo-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran

(S)-3-*N,N*-Dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzo-1-pyran (6.9 g, 16 mmol) and sodium acetate (1.5 g, 18 mmol) were dissolved in acetic acid (430
10 mL). To the solution was added iodine monochloride (18 mL, 1 M, 18 mmol) and the reaction mixture was stirred at room temperature, protected from light, for 24 h. Additional iodine monochloride (2.5 mL, 1M, 2.5 mmol) was added followed by stirring for 3 h. The solvent was evaporated *in vacuo* and the residue was partitioned between methylene chloride (800 mL) and 2 M NaOH (120 mL). The aqueous phase was extracted with
15 methylene chloride (100 mL) and the combined organic layers were washed with brine (2 x 100 mL) and dried (MgSO₄). Evaporation of the solvent gave 8.6 g of a crude product. Purification by column chromatography on silica using ethyl acetate/ethanol (saturated with ammonia) (25:1) as the eluent gave 4.1 g (43% yield) of the title compound (containing about 7% of the starting material) as a yellowish solid: EIMS (70 eV) *m/z*
20 (relative intensity) 553 (15, M⁺). The product was used in the next step without further attempts to purification.

Preparation 40

25 **(S)-8-Carboxymethyl-3-*N,N*-dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran**

(S)-3-*N,N*-Dibenzylamino-8-iodo-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran (2.6 g, 4.8 mmol) was dissolved in *N,N*-dimethylformamide (100 mL) and flushed with carbonmonoxide. To the solution was added palladium acetate (110 mg, 0.48 mmol), 1,3-bis(diphenylphosphino)propane (200 mg, 0.48 mmol), methanol (25 mL) and
30 triethylamine (3.3 mL, 24 mmol). The mixture was reacted with carbonmonoxide at 90 °C and at atmospheric pressure for 8 h. The solution was filtered, the solvent was evaporated.

The residue was co-evaporated with xylene (2 x 50 mL) and partitioned between methylene chloride (300 mL) and 2 M NH₃ (50 mL). The aqueous phase was extracted with methylene chloride (50 mL) and the combined organic layers were washed with brine (2 x 50 mL) dried (MgSO₄). The solvent was evaporated giving 4.0 g of a crude product.

5 Purification by column chromatography on silica using methylene chloride/ethanol (saturated with ammonia) (50:1) as the eluent gave 1.7 g (68% yield) of the title compound (containing about 5% of the corresponding 8-H analogue) as a yellowish solid: EIMS (70 eV) *m/z* (relative intensity) 485 (8, M⁺). The product was used in the next step without further attempts to purification.

10

Preparation 41

(S)-3-*N,N*-Dibenzylamino-8-hydroxymethyl-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran

(S)-8-Carboxymethyl-3-*N,N*-dibenzylamino-5-(methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (490 mg, 1.0 mmol) was dissolved in dry tetrahydrofuran (40 mL) and lithium aluminium hydride (76 mg, 2.0 mmol) was added portionwise. The reaction mixture was stirred at 45 °C for 4 h and cooled to room temperature. The reaction was quenched by the addition of water (76 µL), 15% NaOH (76 µL) and water (225 µL) and stirred for 18h. The white precipitate was filtered off and the solution was dried (MgSO₄). The solvent was
15 evaporated *in vacuo* giving 520 mg of a crude product. Purification by column chromatography on silica using chloroform/ethanol (saturated with ammonia) (15:1) as the eluent gave 390 mg (85% yield) of the title compound containing about 8% of the corresponding 8-methyl analogue) as a yellowish oil: EIMS (70 eV) *m/z* (relative intensity) 457 (15, M⁺).

25

Preparation 42

(S)-3-Amino-8-methyl-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran

(S)-3-*N,N*-Dibenzylamino-8-hydroxymethyl-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (420 mg, 0.90 mmol) was dissolved in methanol (60 mL) and ammonium formate (460 mg, 7.3 mmol) was added. The solution was flushed with nitrogen and palladium on charcoal (120 mg, 10%) was added. The reaction mixture was stirred at 50
30

°C for 16 h. The catalyst was filtered off and the solvent was evaporated *in vacuo* giving 260 mg of a crude product. The residue was dissolved in acetic acid (50 mL) and palladium on charcoal (120 mg, 10%) was added. The reaction mixture was hydrogenated at room temperature and at atmospheric pressure for 46 h. The catalyst was filtered off and the solvent was evaporated *in vacuo*. The residue was partitioned between ethyl acetate (120 mL) and 2 M NaOH (10 mL) and the aqueous phase was extracted with ethyl acetate (10 mL). The combined organic layers were washed with brine (5 mL), dried (MgSO₄) and the solvent was evaporated *in vacuo* giving 200 mg of a crude product. Purification by preparative TLC on silica using chloroform/ethanol (saturated with ammonia) (10:1) as the eluent afforded 150 mg (64% yield) of the title compound as an oil: EIMS (70 eV) *m/z* (relative intensity) 261 (100, M⁺).

Preparation 43

8-Methoxy-5-nitro-3,4-dihydro-2*H*-1-benzopyran-3-carboxylic acid ethyl ester.

To a stirred solution of 8-methoxy-3,4-dihydro-2*H*-1-benzopyran-3-carboxylic acid ethyl ester (5.5 g, 23 mmol; described in: Thorberg, S-O et al. *Acta Pharm.Suec.*1987, 24, (4), 169-182) in methylene chloride (50 mL) at 0 °C was added dropwise 65% HNO₃ (2.0 mL). The solution was stirred at room temperature for 2 h and washed with water. The organic phase was dried and the solvent evaporated *in vacuo*. The residue was treated with diisopropyl ether (30 mL) and ethyl acetate (5 mL) to yield 1.5 g (5.3 mmol) of crystals of the 6-nitro isomer. The mother liquor was purified by column chromatography using diisopropylether as the eluent affording 1.3 g (20% yield) of the title compound: mp 66-68 °C; EIMS (70 eV) *m/z* (relative intensity) 281 (100, M⁺).

Preparation 44

8-Methoxy-5-nitro-3,4-dihydro-2*H*-1-benzopyran-3-carboxylic acid.

A mixture of 8-methoxy-5-nitro-3,4-dihydro-2*H*-1-benzopyran-3-carboxylic acid ethyl ester (5.8 g, 21 mmol) in ethanol (150 mL) and 2 M NaOH (15 mL) was heated to reflux for 30 min. The solvent was evaporated *in vacuo* the residue dissolved in water.

Acidification to pH 2 and extraction with ethyl acetate followed by evaporation of the

solvent *in vacuo* gave 4.9 g (94 % yield) of the title compound: mp 181-183 °C; EIMS (70 eV) *m/z* (relative intensity) 253 (55, M⁺).

Preparation 45

5 ***N*-[4-(4-Morpholinyl)phenyl]-8-methoxy-5-nitro-3,4-dihydro-2*H*-1-benzopyran-3-carboxamide.**

To a solution of 8-methoxy-5-nitro-3,4-dihydro-2*H*-1-benzopyran-3-carboxylic acid (2.5 g, 10 mmol) in toluene (40 mL) and *N,N*-dimethylformamide (1 mL) was added thionyl chloride (3.6 mL, 50 mmol). The reaction mixture was refluxed for 2 h and the solvent was removed *in vacuo*. The residual acid chloride was added to a solution of 4-(1-morpholino)aniline (1.78 g, 10 mmol; described in: Devlin, J.P. et. al., *J. Chem. Soc. Perkin Trans, 1*. 1975 830-841) and triethylamine (2.0 g, 20 mmol) in methylene chloride (30 mL) and stirred at 0 °C for 10 min and for 1 h at room temperature. The solvent was removed *in vacuo* and the residue was dissolved in ethyl acetate and washed with 2 M NaOH. Evaporation of the solvent *in vacuo* afforded 1.5 g (36 % yield) of the title compound as white crystals: mp 238-240 °C; EIMS (70 eV) *m/z* (relative intensity) 413 (5, M⁺).

Preparation 46

20 ***N*-[4-(4-Morpholinyl)phenyl]-5-amino-8-methoxy-3,4-dihydro-2*H*-1-benzopyran-3-carboxamide.**

To a solution of *N*-[4-(4-morpholinyl)phenyl]-8-methoxy-5-nitro-3,4-dihydro-2*H*-1-benzopyran-3-carboxamide (1.2 g, 2.9 mmol) in *N,N*-dimethylformamide (10 mL) was added a solution of sodium dithionite (2.1 g, 12 mmol) in water (5 mL). The mixture was stirred at 55 °C for 3 h and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica gel using ethyl acetate as the eluent affording 273 mg of the title compound (55% yield): EIMS (70 eV) *m/z* (relative intensity) 383 (100, M⁺).

Preparation of the 5-HT_{1B} antagonist compounds described herein

Example 1

(R)-N-[8-(Piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

To an ice-cooled solution of (R)-N-[8-(4-*tert*-butoxycarbonylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinophenylcarboxamide (1.0 g, 2 mmol) in methylene chloride (100 mL) was added trifluoroacetic acid (3 mL). The reaction was stirred at ambient temperature for 7 h. The solvent was evaporated *in vacuo* and the residue was dissolved in water (20 mL), alkalized with a 2 M aqueous sodium hydroxide solution and extracted, twice, with methylene chloride. The phases were separated, the combined organic phases were dried (Na₂SO₄), filtered and evaporated *in vacuo*. Purification on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (95:5:0.5) as the eluent gave 580 mg (70% yield) of the title compound as white crystals: mp 202-203 °C; [α]_D²¹ -56° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 420 (5, M⁺).

Example 2

(R)-N-[8-(4-Ethylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

To a solution of (R)-N-8-(piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide (90 mg, 0.21 mmol) in acetone (20 mL) were added potassium carbonate (44 mg, 0.32 mmol) and iodoethane (26 μ L, 0.32 mmol) and the reaction was stirred for 48 h at ambient temperature. The reaction mixture was filtered and the solvent evaporated *in vacuo*. The residue was partitioned between methylene chloride and water, the phases were separated, and the organic phase was dried (Na₂SO₄), filtered and evaporated *in vacuo*. Purification on a silica gel column using chloroform/ethanol saturated with NH₃ (100:3) as the eluent gave 63 mg (66% yield) of the title compound as white crystals: mp: 204-206 °C; [α]_D²¹ -67° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 448 (21, M⁺).

Example 3**(R)-N-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide**

To a solution of 4-morpholinobenzoic acid (0.92 g, 4.5 mmol; described in: Degutis, J.; Rasteikiene, L.; Degutiene, A. *Zh. Org. Khim.* **1978**, *14*(10), 2060-2064) in anhydrous *N,N*-dimethylformamide (75 mL) was added 1,1'-carbonyldiimidazole (0.76 g, 4.8 mmol) and the reaction was heated at 75 °C. When the carbon dioxide evolution had ceased (after 45 min), the reaction was cooled to room temperature and a solution of (*R*)-2-amino-5-methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (1.2 g, 4.2 mmol) dissolved in anhydrous *N,N*-dimethylformamide (20 mL) was added. The reaction was allowed to stir at ambient temperature for 48 h and the solvent was evaporated *in vacuo*. Purification on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (180:5:0.5) as the eluent followed by recrystallization from ethyl acetate and a few drops of methanol gave 1.0 g (53% yield) of white crystals: mp 237-238 °C [α]_D²¹ - 40 ° (c=1, chloroform); EIMS (70eV) *m/z* (relative intensity) 464 (5, M⁺).

Example 4**(R)-N-[5-Ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide**

4-Morpholinobenzoic acid (64 mg, 0.31 mmol) was dissolved in dry *N,N*-dimethylformamide (1 mL) and 1,1'-carbonyldiimidazole (52 mg, 0.32 mmol) was added. The reaction mixture was stirred at 75 °C for 1 h and cooled to room temperature. A solution of (*R*)-2-amino-5-ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (80 mg, 0.29 mmol) in dry *N,N*-dimethylformamide (3 mL) was added and the reaction mixture was stirred at room temperature for 14 h. The solvent was evaporated and the residue was dried *in vacuo*. The crude product was purified by preparative TLC on silica using chloroform/methanol/conc. NH₃ (95:5:0.5) as the eluent which gave 85 mg (59% yield) of the title compound as a white solid: mp 234 °C (dec); EIMS (70 eV) *m/z* (relative intensity) 462 (27, M⁺); [α]_D²¹ -48° (c 0.09, chloroform).

Example 5**(R)-N-[5-Ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-(4-morpholinocarbonyl)benzamide**

4-Morpholinocarbonylbenzoic acid (180 mg, 0.77 mmol; described in: *J. Med. Chem.* 1994, 37(26), 4538-4554) and 1,1'-carbonyldiimidazole (130 mg, 0.80 mmol) were dissolved in dry *N,N*-dimethylformamide (3 mL) and stirred at 75 °C for 2 h. After cooling to room temperature, a solution of (*R*)-2-amino-5-ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (200 mg, 0.73 mmol) in dry *N,N*-dimethylformamide was added and the reaction mixture was stirred for 60 h. The solvent was evaporated *in vacuo* and the residue was partitioned between methylene chloride (60 mL) and 2 M NH₃ (5 mL). The organic phase was washed with brine (10 mL) and dried (Na₂SO₄). Evaporation of the solvent *in vacuo* gave 360 mg of a crude product. Purification by column chromatography on silica using chloroform/methanol/conc. NH₃ (95:5:0.5) as the eluent afforded 240 mg (65% yield) of the title compound as a white solid: mp 213-214 °C; EIMS (70 eV) *m/z* (relative intensity) 490 (27, M⁺); [α]²¹_D -28° (c 0.15, chloroform).

Example 6**(R)-N-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinocarbonylbenzamide**

The title compound was prepared from (*R*)-2-amino-5-methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene following the general method of Preparation 16. Purification on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (96:4:0.3) as the eluent gave after recrystallization from ethyl acetate/diethyl ether 93 mg (52% yield) of white crystals: mp 209-210 °C; [α]²¹_D -18° (c=1, chloroform); EIMS (70eV) *m/z* (relative intensity) 492 (36, M⁺).

Example 7**(R)-N-[5-Bromo-8-(piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide**

To an ice-cooled solution of (*R*)-N-[5-bromo-8-(4-*tert*-butyl oxycarbonylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpho-

linobenzamide (150 mg, 0.26 mmol) in methylene chloride (20 mL) was added trifluoroacetic acid (0.7 mL). The reaction was stirred at ambient temperature for 20 h. The solvent was evaporated *in vacuo* and the residue was dissolved in water (20 mL), alkalized with a 2 M aqueous solution of sodium hydroxide and extracted with methylene chloride.

5 The phases were separated and the organic phase was dried (Na_2SO_4), filtered and evaporated *in vacuo*. The residue was purified on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (90:10:1) as the eluent. Yield: 94 mg (72%) of a white crystals: mp 228-229 °C; $[\alpha]^{21}_{\text{D}} -6^\circ$ ($c=1$, chloroform); EIMS (70 eV) m/z (relative intensity) 498 and 500 (1.5, M^+)

10

Example 8

(R)-N-[5-Bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

The title compound was prepared from (R)-2-amino-5-bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene following the general method of Preparation 16. Purification on a silica gel column using chloroform/methanol/concentrated ammonium hydroxide (95:5:1) as the eluent gave 100 mg (62% yield) of white crystals: mp 245-246 °C $[\alpha]^{21}_{\text{D}} -23^\circ$ ($c=1$, chloroform); EIMS (70eV) m/z (relative intensity) 512 and 514 (1, M^+).

15

Example 9

(R)-N-[5-Bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-trifluoromethylbenzamide

(R)-N-[8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-trifluoromethylbenzamide (80 mg, 0.19 mmol) and sodium acetate (200 mg) were dissolved in acetic acid (3 mL) and the mixture was stirred at room temperature. Bromine (34 mg, 0.21 mmol) was added dropwise to the reaction mixture and the mixture was stirred for 2 h at ambient temperature. A 2 M sodium hydroxide solution (100 mL) was added and the mixture was extracted with diethyl ether (2x50 mL). The combined organic phases were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*.

25

30 Purification on a silica gel column using methylene chloride/ethanol saturated with NH_3

(94:6) as the eluent gave 80 mg (85% yield) of the title compound as a white solid: mp 229-230 °C; $[\alpha]_D^{21}$ -5.4° ($c=1$, chloroform); EIMS (70 eV) m/z (relative intensity) 495 and 497 (3, M^+).

5 **Example 10**

(R)-N-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

4-Morpholinobenzoic acid (92 mg, 0.44 mmol) was dissolved in dry *N,N*-dimethylformamide (2 mL) and flushed with nitrogen. To the solution was added 1,1'-carbonyldiimidazole (76 mg, 0.47 mmol) and the reaction mixture was stirred at 75 °C for 1.5 h. The solution was cooled to room temperature and (*R*)-2-amino-5-methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (110 mg, 0.42 mmol), dissolved in dry *N,N*-dimethylformamide (2 mL) was added. The solution was stirred at room temperature for 30 h. The solvent was evaporated *in vacuo* giving 290 mg of a crude product. Purification by preparative TLC on silica gel using chloroform/methanol/conc. NH_3 (95:5:0.5) as the eluent afforded 145 mg (73% yield) of the title compound as a white solid: mp >231 °C (dec); EIMS (70 eV) m/z (relative intensity) 448 (3, M^+); $[\alpha]_D^{21}$ -60° (c 0.15, chloroform).

20 **Example 11**

N-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide

A solution of *N*-(4-morpholinophenyl)-8-amino-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide (1.4 g, 3.5 mmol), bis (2-chloroethyl)-methylamine hydrochloride (960 mg, 5 mmol) and sodium hydrogen carbonate (420 mg, 5 mmol) in *n*-butanol (30 mL) was heated at 90 °C for 5 h. After cooling, 2 M ammonium hydroxide (30 mL) was added and the mixture heated at 50 °C for 1 h. The phases were separated, evaporated *in vacuo* and purified by flash chromatography on a silica gel column with chloroform/ethanol/conc. ammonium hydroxide 90/10/0.3 as eluent. Yield: 320 mg (20 %) of the title compound: mp 230-232 °C; EIMS (70eV) m/z (relative intensity) 464 (75, M^+).

Chromatographic Preparation of the Enantiomers of *N*-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide

N-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide (5 mg) was dissolved in 4 ml of eluent consisting of acetonitrile and pH 3.0 phosphate buffer, $\mu = 0.1$ (62.5 : 37.5, v/v). This solution was purified on a Nucleosil 7 C₁₈ column (25 x 250 mm) with the above mobile phase to remove late eluting impurities. The collected fractions of the main component were concentrated under reduced pressure at 35-39 °C. The residue was dissolved in 30 ml of the eluent composed of 10 mM ammonium acetate, diethylamine and acetic acid (4000+2+2, v/v/v, pH 5.26) and the chiral semi-preparation of the enantiomers of *N*-(4-morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide was carried out on a Chiral AGP semi-preparative column (10 x 150 mm) using a guard column of the same stationary phase. 2.0 ml/min of flow rate was used and detection was monitored at 260 nm. Fractions of both enantiomers were separately collected and concentrated to a volume of about 5 ml under reduced pressure at 35-39 °C. The concentrated fractions were adjusted to pH 10-11 with 5 M NaOH and extracted with chloroform. The two organic phases were washed with water and dried with anhydrous magnesium sulfate. After being filtered through glasswool, the organic filtrates were evaporated *in vacuo* affording the two enantiomers as two slightly yellow solids.

Example 12

***N*-(Morpholinocarbonylphenyl)-8-(4-methylpiperazin-1-yl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide**

A solution of *N*-(morpholinocarbonylphenyl)-8-amino-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide (280 mg, 0.69 mmol), bis (2-chloroethyl)methylamine hydrochloride (190 mg, 1.0 mmol) and sodium hydrogen carbonate (84 mg, 1.0 mmol) in *n*-butanol (20 mL) was heated at 90 °C for 5 h. After cooling, 2 M ammonium hydroxide (10 mL) was added and the mixture was heated at 50 °C for 1 h. The organic phase was evaporated *in vacuo* and the residue was purified by flash chromatography on a

silica gel column using chloroform/ethanol/conc. ammonium hydroxide (90:10:0.5) as eluent to yield 60 mg (18%) of the title compound: EIMS (70eV) m/z (relative intensity) 492 (50, M^+).

5 **Example 13**

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-morpholinobenzamide.

A solution of 4-morpholinobenzoic acid (380 mg, 1.83 mmol, described in: Degutis, J.; Rasteikiene, L.; Degutiene, A. *Zh. Org. Khim.* **1978**, *14*(10), 2060-2064) and 1,1'-
10 carbonyldiimidazole (310 mg, 1.92 mmol) in anhydrous *N,N*-dimethylformamide (12 mL) was stirred at 75 °C for 30 min. The mixture was allowed to cool after which a solution of (*S*)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran (430 mg, 1.74 mmol) in *N,N*-dimethylformamide (8 mL) was added. The reaction mixture was stirred at room temperature for 3 days. Another portion of 1,1'-carbonyldiimidazole (57 mg, 0.35
15 mmol) was added, and the mixture was stirred for an additional 3.5 h. The solvent was removed *in vacuo*, and the residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 93:7 + 0.5% NH_3) affording 513 mg (68% yield) of the title compound as a white solid: mp 210-212 °C; $[\alpha]^{22}_D$ -145° (*c* 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 436 (65, M^+).

20

Example 14

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(4-piperidon-1-yl)benzamide.

A solution of 1,1'-carbonyldiimidazole (116 mg, 0.716 mmol) and 4-(4-piperidon-1-yl)benzoic acid (150 mg, 0.683 mmol) in anhydrous *N,N*-dimethylformamide (5 mL) was
25 stirred at 75 °C for 50 min. The mixture was allowed to cool, and a solution of (*S*)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran (161 mg, 0.651 mmol) in *N,N*-dimethylformamide (4 mL) was added. The reaction mixture was stirred at room temperature for 8 days. The solvent was removed *in vacuo*, and the residue was purified by
30 column chromatography on silica (eluent: chloroform/ethanol, 90:10 + 0.5% conc. NH_3)

affording 54 mg (19% yield) of the title compound as a white solid: mp 222-225 °C (decomposes); $[\alpha]_D^{22}$ -136° (*c* 0.30, chloroform); TSPMS (70eV) *m/z* 449 (M+1).

Example 15

5 (S)-N-[8-Methyl-5-(4-methyl-piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(dimethylaminocarbonyl)benzamide

4-(Dimethylaminocarbonyl)benzoic acid /Jurewicz, A.T; U.S Patent 3,607,918 1971) (38 mg 0.20 mmol) and 1,1'-carbonyldiimidazole (34 mg, 0.21 mmol) were dissolved in dry *N,N*-dimethylformamide (4 mL) and stirred at 75° C for 1.5 h. The reaction mixture was cooled to room temperature and a solution of (S)-3-amino-8-methyl-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (49 mg 0.19 mmol) in dry *N,N*-dimethylformamide (5mL) was added. The reaction mixture was stirred at 50° C for 14 h and the solvent was evaporated *in vacuo* giving 120 mg of crude product. Purification by preparative TLC using chloroform/ methanol/conc.NH₃ (95:5:0.5) as the eluent afforded 40 mg (48% yield) of the title compound as a white foam: EIMS (70 eV) *m/z* (relative intensity) 436 (26, M⁺); $[\alpha]_D^{21}$ -9° (*c* 0.20, chloroform).

Example 16

20 N-[4-(4-Morpholinyl)phenyl]-8-methoxy-5-(4-methyl-piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-carboxamide.

A solution of N-[4-(4-morpholinyl)phenyl]-5-amino-8-methoxy-3,4-dihydro-2H-1-benzopyran-3-carboxamide (270 mg, 0.7 mmol), bis (2-chloroethyl)-methylamine hydrochloride (288 mg, 1.5 mmol) and sodium hydrogen carbonate (126 mg, 1.5 mmol) in *n*-butanol (10 mL) was stirred at 90 °C for 2.5 h. 2 M ammonia (10 mL) was added at 50 °C, the mixture was cooled and the phases were separated. The organic phase evaporated *in vacuo* and the residue was purified by column chromatography on silica gel using ethyl acetate/triethyl amine (100:8) as the eluent affording 170 mg (50% yield) of the title compound as white crystals: mp 202-204 °C; EIMS (70 eV) *m/z* (relative intensity) 466 (100 M⁺).

Example 17**(R)-N-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide**

To a solution of 4-morpholinobenzoic acid (0.89 g, 4.3 mmol; described in: Degutis, J.;
5 Rasteikiene, L.; Degutiene, A. *Zh. Org. Khim.* **1978**, *14*(10), 2060-2064) in anhydrous
N,N-dimethylformamide (30 mL) was added 1,1'-carbonyldiimidazole (0.73 g, 4.3 mmol)
and the reaction was heated at 75 °C. When the carbon dioxide evolution had ceased (after
30 min), the reaction was cooled to room temperature and a solution of (*R*)-2-amino-8-(4-
10 methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthalene (1.0 g, 4.1 mmol) in anhydrous *N,N*-
dimethylformamide (5 mL) was added. The reaction was allowed to stir at ambient
temperature for 24 h and the solvent was evaporated *in vacuo*. Purification on a silica gel
column using chloroform/methanol/ concentrated ammonium hydroxide (95:5:0.5) as the
eluent gave 1.5 g (85% yield) of the title compound as white crystals: mp 230-231 °C;
[α]_D²¹ -49° (*c* 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 434 (10, M⁺).

Pharmacology**Methods for testing****(i) Functional h5-HT_{1B} receptor assay**

In order to evaluate the antagonistic properties of the 5-HT_{1B}-receptor the standard assay
using electrical field stimulation of [³H] -5-HT release from occipital cortex of guinea pigs
can be used.

Methods and Materials:

30 **Buffer composition (mM)** NaHCO₃ (25), NaH₂PO₄. H₂O (1.2), NaCl (117), KCl(6),
MgSO₄·7H₂O(1.2), CaCl₂(1.3), EDTA Na₂(0.03). The buffer is gassed for at least 30 min

before use. The pH of the buffer is about 7.2 of room temperature but it rises to about 7.4 at 37°C.

Preparation of occipital cortical slices

- 5 Guinea pigs (200-250 g) were decapitated and the whole brains were removed. The occipital cortices were dissected and cut into slices 0.4x4 mm with a McIlwain chopper machine. The white part of the tissue was removed carefully with a tweezer before slicing. The slices were incubated in 5 ml buffer in the presence of 5 mM pargyline chloride. After incubation with 0.1 mM [³H]-5-HT for another 30 min the slices were transferred to a test
10 tube and washed three times with the same volume of buffer. The slices were transferred to the superfusion chambers with a plastic pipette and were washed for 40 min. with the buffer in the presence of uptake inhibitor citalopram (2.5 µM) with a flow of 0.5 ml/min.

Electrical stimulation of 5-HT release

- 15 The superfused buffer was collected in 2 ml fractions. The slices were stimulated by electricity with a train of pulses of frequency 3 Hz, duration 2 ms and current 30 mA for 3 min at the 4th and 13th fractions. The tested drugs were added from the 8th fraction to the end of the experiment.

Results

- 20 A first electrical (or K⁺) stimulation resulted in a standard amount of [³H] 5-HT released (S₁). Between the first and second stimulation the h5-HT_{1B} antagonist is added to the media which results in a dose depending increase of the release(S₂) during the second stimulation. The S₂/S₁ ratio which is the per cent of released [³H] 5-HT at the second
25 stimulation (S₂) divided by that of the first stimulation (S₁) was used to estimate drug effects on transmitter release. See Fig.1.

(ii) effects of h5-HT_{1B} receptor antagonists/partial agonists in combination with SSRIs measured by microdialysis in guinea pig brain

5 The effects of different h5-HT_{1B} receptor antagonists in combination with a selective serotonin reuptake inhibitor on extracellular levels of 5-HT in the guinea pig frontal cortex is measured by microdialysis.

Male Dunkin Hartley guinea pigs (Charles River, Sweden), weighing 270-500 g, were
10 anaesthetized intramuscularly with a 1:3 v/v mixture of Rompun® vet (20 mg/ml) and Ketalar® vet (50 mg/ml), and placed in a stereotaxic frame. A unilateral guide cannula was carefully implanted into the frontal cortex using the following stereotaxic coordinates with respect to bregma: AP: +4.5 mm, L: -2.0 mm and DV: 0 mm from the brain surface. The animals were allowed to recover for 2-7 days before the experiment were carried out.
15 The day before starting dialysis sampling, microdialysis probes with 3 mm membranes were inserted into the guide cannula. The probes were perfused with Ringer solution at a flow rate of 2.0 µl/min and samples were collected every 20 min. Compound A (a selective serotonin reuptake inhibitor, SSRI) was given at time 0 min and compound B, C, D or E (a h5-HT_{1B} antagonist or saline) was administered 60 min later. All drugs were given
20 subcutaneously. The 5-HT content was analyzed by high performance liquid chromatography (HPLC) with electrochemical detection. For data analysis, the average 5-HT value of 3-4 samples collected pre-administration of drugs was defined as 100% (baseline) and the following samples expressed as percent of this value.

25 **Results**

Fig.2. shows the effects of compound A in combination with compound B, C, D or E on extracellular levels of 5-HT in the guinea pig frontal cortex as measured by microdialysis. Data are mean ± S.E.M; n=5-6. The arrows indicate drug administration.

Compound A: Citalopram

Compound B: (*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

5

Compound C: Saline

Compound D: (*R*)-*N*-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide


10

Compound E: (*R*)-*N*-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide

15

Claims.

1. A combination of a first component (a) which is a 5-HT reuptake inhibitor with a
 second component (b) which is a selective h5-HT_{1B} antagonist or partial agonist having
 the formula I



(I)


wherein X is CH₂, O;

Y is CONH, NHCO;

R₁ is H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl;

R₂ is H, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen;

R₃ is



R₄ and R₅ independently are H or C₁-C₄ alkyl,

as racemate, *R*-enantiomer or *S*-enantiomer, and said components (a) and (b) being in the
 form of free bases, solvates or pharmaceutically acceptable salts thereof.

2. A combination according to claim 1 wherein the second component (b) is a compound of formula I wherein X is CH₂.

3. A combination according to claim 2 wherein the second component (b) is a compound
5 of formula I wherein Y is NHCO.

4. A combination according to claim 3 wherein the second component (b) is a compound of formula I wherein R₃ is morpholino.

10 5. A combination according to any one of claims 1-4 wherein the second component (b) is a compound of formula I wherein R₁ is hydrogen, methyl or ethyl and R₂ is hydrogen, methyl, ethyl, methoxy or bromo.

6. A combination according to claim 1 wherein the second component (b) is a compound
15 selected from

(*R*)-*N*-[8-(Piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[8-(4-Ethylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

20 (*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[5-Ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

25 (*R*)-*N*-[5-Ethyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinocarbonyl benzamide;

(*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinocarbonylbenzamide;

(*R*)-*N*-[5-Bromo-8-(piperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

N-[5-Bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

(*R*)-*N*-[5-Bromo-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-trifluoromethylbenzamide;

5 (*R*)-*N*-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide;

N-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

10 (*R*)-*N*-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

(*S*)-*N*-(4-Morpholinophenyl)-8-(4-methylpiperazinyl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

(*R*)-*N*-(Morpholinocarbonylphenyl)-8-(4-methylpiperazin-1-yl)-5-methoxy-1,2,3,4-tetrahydronaphthalene-2-carboxamide;

15 (*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-morpholinobenzamide;

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(4-piperidon-1-yl)benzamide;

20 (*S*)-*N*-[8-Methyl-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(dimethylaminocarbonyl)benzamide;

N-[4-(4-Morpholinyl)phenyl]-8-methoxy-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-carboxamide,

7. A combination according to claim 6 wherein the second component (b) is a compound
25 selected from (*R*)-*N*-[8-(4-Methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide, (*R*)-*N*-[5-Methoxy-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide and (*R*)-*N*-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydro-2-naphthyl]-4-morpholinobenzamide.

30 8. The combination according any one of claims 1-7 wherein the

5-HT reuptake inhibitor is fluoxetine, paroxetine, citalopram, clomipramine, sertraline or fluvoxamin.

9. Use of the combination according to any one of claims 1-8 for the manufacture of a medicament for the treatment of affective disorders.

10. Use of the combination according to claim 9 for the manufacture of a medicament for the treatment of depression.

11. A method for the treatment of affective disorders by administering to a patient suffering therefrom the combination defined in any one of claims 1-8.

12. A method for the treatment of depression by administering to a patient suffering therefrom the combination defined in any one of claims 1-8.

13. A pharmaceutical formulation wherein the active ingredients are the combination defined in any one of claims 1-8 optionally in association with adjuvants, diluents, excipients and/or inert carriers.

14. A pharmaceutical formulation according to claim 13 wherein the first component (a) is concomitantly administered with the second component (b).

15. A pharmaceutical formulation according to any one of claims 13-14 for use in the treatment of affective disorders.

16. A pharmaceutical formulation according to claim 15 for use in the treatment of depression.

17. A process for the preparation of the combination according to claim 1 whereby a 5-HT reuptake inhibitor as defined in any one of claims 1 or 8 is incorporated into the same

pharmaceutical formulation as a selective 5-HT_{1B} antagonist or partial agonist as defined in any one of claims 1-7.


18. A process for the preparation of the combination according to claim 1 whereby a 5-HT reuptake inhibitor as defined in any one of claims 1 or 8 is incorporated into the same pharmaceutical formulation and a selective 5-HT_{1B} antagonist or partial agonist as defined in any one of claims 1-7 is in a pharmaceutical formulation.

19. A kit containing the combination of a first component (a) which is a 5-HT reuptake inhibitor as defined in any one of claims 1 or 8 and a second component (b) which is selective 5-HT_{1B} antagonist or partial agonist as defined in any one of claims 1-7, optionally with instructions for use.

20. A method of improving the onset of therapeutic action by concomitant administration of a first component (a) which is a 5-HT reuptake inhibitor as defined in any one of claims 1 or 8 and a second component (b) which is selective 5-HT_{1B} antagonist or partial agonist as defined in any one of claims 1-7.


1 / 2

(R)-N-[5-Methyl-8-(4-methylpiperazin-1-yl)-1,2,3,4-tetrahydronaphthyl]-4-morpholinobenzamide



2 / 2

Fig. 2



INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/01601

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A61K 31/495, A61K 31/535

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Brain Research Bulletin, Volume 43, No 4, 1997, Colin Davidson et al, "Synergism of 5-HT 1B/D Antagonists with Paroxetine on Serotonin Efflux in Rat Ventral Lateral Geniculate Nucleus Slices" page 405 - page 409 --	1-10,13-19
X	EP 0533267 A1 (GLAXO GROUP LIMITED), 24 March 1993 (24.03.93), page 5, line 26 - line 35, claim 16 --	1-10,13-19
X	EP 0533268 A1 (GLAXO GROUP LIMITED), 24 March 1993 (24.03.93), page 6, line 44 - line 53, claim 19 --	1-10,13-19

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

27 January 1999

Date of mailing of the international search report

28 -01- 1999

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Solveig Gustavsson

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/01601

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	WO 9734883 A1 (ASTRA AKTIEBOLAG), 25 Sept 1997 (25.09.97) -- -----	1-10,13-19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/01601

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11-12, 20
because they relate to subject matter not required to be searched by this Authority, namely:

See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

21/12/98

International application No.

PCT/SE 98/01601

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
EP	0533267	A1	24/03/93	AU	2452892 A	25/03/93
				AU	2568792 A	27/04/93
				CA	2078507 A	19/03/93
				CN	1073430 A	23/06/93
				CZ	9400611 A	16/11/94
				FI	941261 A	17/03/94
				JP	6107637 A	19/04/94
				MX	9205278 A	01/03/93
				NO	940974 A	17/03/94
				US	5358948 A	25/10/94
				WO	9306084 A	01/04/93
				ZA	9207106 A	17/03/94
EP	0533268	A1	24/03/93	AP	303 A	28/01/94
				AP	9200427 D	00/00/00
				AU	656021 B	19/01/95
				AU	2453092 A	25/03/93
				CA	2078505 A	19/03/93
				CN	1071919 A	12/05/93
				FI	924160 A	19/03/93
				HU	65608 A	28/07/94
				IL	103198 A	18/06/96
				JP	6116251 A	26/04/94
				MX	9205280 A	01/03/93
				NZ	244373 A	28/03/95
				RU	2077535 C	20/04/97
				US	5340810 A	23/08/94
				US	5510350 A	23/04/96
				ZA	9207108 A	08/09/93
				CN	1076195 A	15/09/93
WO	9734883	A1	25/09/97	AU	2186597 A	10/10/97
				AU	6949796 A	27/03/97
				HR	970166 A	30/04/98
				SE	9601110 D	00/00/00